

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 February 2003 (20.02.2003)


PCT

(10) International Publication Number  
WO 03/013430 A2


- (51) International Patent Classification<sup>7</sup>: **A61K**
- (21) International Application Number: PCT/US02/24891
- (22) International Filing Date: 5 August 2002 (05.08.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/310,079 6 August 2001 (06.08.2001) US  
60/389,255 14 June 2002 (14.06.2002) US  
60/393,929 3 July 2002 (03.07.2002) US  
Not furnished 12 July 2002 (12.07.2002) US
- (71) Applicant: **KOSAN BIOSCIENCES, INC.** [US/US];  
3832 Bay Center Place, Hayward, CA 94545 (US).
- (72) Inventors: **SANTI, Daniel**; 211 Belgrave Avenue, San Francisco, CA 94111 (US). **MYLES, David, C.**; 1 Eagle Hill, Kensington, CA 94707 (US). **TIAN, Zq**; 5029 Xavier Common, Fremont, CA 94555 (US). **HUTCHINSON, C., Richard**; 423 Highland Avenue, San Mateo, CA 94401 (US). **JOHNSON, Robert**; 3656 Happy Valley Road, Lafayette, CA 94549 (US). **ZHOU, Yi-Qing**; 1153 Camino Vallecito, Lafayette, CA 94549 (US). **FENG, Li**; 5351 Laramie Court, Fremont, CA 94536 (US).
- (74) Agents: **FAVORITO, Carolyn, A.** et al.; Morrison & Forster LLP, Suite 500, 3811 Valley Centre Drive, San Diego, CA 92130-2332 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,

[Continued on next page]

(54) Title: BENZOQUINONE ANSAMYCINS




geldanamycin



17-allylamino-17-desmethoxy-  
geldanamycin  
(17-AAG)

(57) Abstract: The invention relates to benzoquinone ansamycin analogs useful for the treatment of cancer and other diseases or conditions characterized by undesired cellular proliferation or hyperproliferation. Therapies involving the administration of such benzoquinone ansamycin analogs, optionally in combination with an inhibitor of an HSP90 client protein, are useful to treat cancer and non-cancerous disease conditions.



17-(2-dimethylaminoethyl)amino-17-  
desmethoxygeldanamycin  
(17-DMAG)



MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

**Published:**

— without international search report and to be republished upon receipt of that report

- (84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## BENZOQUINONE ANSAMYCINS

### CROSS-REFERENCE TO RELATED APPLICATIONS

5 [0001] This application claims priority to U.S. Provisional Application  
60/310,079, filed August 6, 2001, entitled "Geldanamycin Analogs," and to U.S.  
Provisional Applications 60/389,255, filed 14 June 2002; 60/393,929, filed July 3,  
2002 (Atty. Docket No. 30097.01); and 60/\_\_\_\_, filed July 12, 2002 (Atty.  
Docket No. 30097.02); each entitled "Recombinant Geldanamycin Polyketide  
Synthase Genes." Each of the above documents is incorporated herein by reference  
10 in their entireties.

### STATEMENT OF GOVERNMENT INTEREST

15 [0002] The present invention was made in part under NIH Grant No. R43  
CA96262-01. The United States government may have certain rights in this  
invention.

### BACKGROUND

20 [0003] The clinical utility of many potential anti-cancer compounds is  
limited by undesired toxicity against non-target cells. Undesired toxicity in a drug is  
typically due to a lack of specificity, either in target tissue or in mechanism of  
action. If the drug target is present in normal as well as diseased tissues, then  
normal tissue as well as diseased tissue may be affected by the drug. The drug may  
25 also have multiple mechanisms of toxicity, one specific for diseased cells and the  
other non-specific. In either instance, there is often a dose-dependent difference in  
the actions of drugs against normal and diseased tissues, with the effects on diseased  
tissues being observed at lower concentrations than the effects on normal tissues. A  
major task of anti-cancer therapy is thus to determine the dosage at which the drug is  
30 therapeutically effective with minimal effects on normal tissues.

[0004] Phase I clinical trials are typically used to determine the maximum  
tolerated dose (MTD) of a potential anti-cancer compound, i.e., the maximum dose

that can be safely administered without incurring toxicity. The difference between the MTD and the therapeutically effective dose is known as the therapeutic window. For a large number of anti-cancer agents, the MTD is very close to the therapeutically effective dose, i.e., the therapeutic window is very small. The MTD may even be lower than the therapeutically effective dose, making the agent unusable in the clinic.

[0005] Clinical anti-cancer therapy often involves attempting to achieve a delicate balance between effectiveness and undesired toxicity. Agents which synergize the action of a drug against diseased tissues while not affecting the toxicity against normal tissues could allow the effective use of doses of drug well below the MTD, thus increasing the therapeutic window and enhancing the safety and effectiveness of the therapy.

[0006] Geldanamycin (Figure 1) is a benzoquinone ansamycin polyketide isolated from *Streptomyces geldanus*. Although originally discovered by screening microbial extracts for antibacterial and antiviral activity, geldanamycin was later found to be cytotoxic to certain tumor cells *in vitro* and to reverse the morphology of cells transformed by the Rous sarcoma virus to a normal state.

[0007] Geldanamycin's nanomolar potency and apparent specificity for aberrant protein kinase dependent tumor cells, as well as the discovery that its primary target in mammalian cells is the ubiquitous Hsp90 protein chaperone, has stimulated interest in the development of this anti-cancer drug. However, the association of hepatotoxicity with the administration of geldanamycin led to its withdrawal from Phase I clinical trials. As with several other promising anticancer agents, geldanamycin also has poor water solubility that makes it difficult to deliver in therapeutically effective doses.

[0008] More recently, attention has focused on 17-amino derivatives of geldanamycin, in particular 17-(allylamino)-17-desmethoxygeldanamycin (17-AAG; Figure 1), that show reduced hepatotoxicity while maintaining Hsp90 binding. Certain 17-amino derivatives of geldanamycin, 11-oxogeldanamycin, and 5,6-dihydrogeldanamycin, are disclosed in U.S. patents 4,261,989; 5,387,584; and 5,932,566, each of which is incorporated herein by reference. Like geldanamycin, 17-AAG has limited aqueous solubility. This property requires the use of a

solubilizing carrier, most commonly Cremophore® (BASF Aktiengesellschaft), a polyethoxylated castor oil which can result in serious side reactions in some patients.

5 [0009] Treatment of cancer cells with geldanamycin or 17-AAG causes a retinoblastoma protein-dependent G1 block, mediated by down-regulation of the induction pathways for cyclin D-cyclin dependent cdk4 and cdk6 protein kinase activity. Cell cycle arrest is followed by differentiation and apoptosis. G1 progression is unaffected by geldanamycin or 17-AAG in cells with mutated retinoblastoma protein; these cells undergo cell cycle arrest after mitosis, again followed by apoptosis.

10 [00010] The mechanism of action of benzoquinone ansamycins appears to be via binding to Hsp90 and subsequent degradation of Hsp90-associated client proteins. Among the most sensitive client protein targets of the benzoquinone ansamycins are the Her kinases (also known as ErbB), Raf, Met tyrosine kinase, and the steroid receptors. Hsp90 is also involved in the cellular response to stress, including heat, radiation, and toxins. Certain benzoquinone ansamycins, such as 17-AAG, have thus been studied to determine their interaction with cytotoxins that do not target Hsp90 client proteins.

15 [00011] U.S. Patents 6,245,759; 6,306,874; and 6,313,138, each of which is incorporated herein by reference, disclose compositions comprising certain tyrosine kinase inhibitors together with 17-AAG and methods for treating cancer with such compositions. Münster *et al.*, "Modulation of Hsp90 function by ansamycins sensitizes breast cancer cells to chemotherapy-induced apoptosis in an RB- and schedule-dependent manner," *Clinical Cancer Research* (2001) 7: 2228-2236, discloses that 17-AAG sensitizes cells in culture to the cytotoxic effects of paclitaxel and doxorubicin. The Münster reference further discloses that the sensitization towards paclitaxel by 17-AAG is schedule-dependent in retinoblastoma protein-producing cells due to the action of these two drugs at different stages of the cell cycle: treatment of cells with a combination of paclitaxel and 17-AAG is reported to give synergistic apoptosis, while pretreatment of cells with 17-AAG followed by treatment with paclitaxel is reported to result in abrogation of apoptosis. Treatment of cells with paclitaxel followed by treatment with 17-AAG 4 hours later is reported to show a synergistic effect similar to coincident treatment.

20

25

30

[00012] Citri *et al.*, "Drug-induced ubiquitylation and degradation of ErbB receptor tyrosine kinases: implications for cancer chemotherapy," *EMBO Journal* (2002) 21: 2407-2417, discloses an additive effect upon co-administration of geldanamycin and an irreversible protein kinase inhibitor, CI-1033, on growth of ErbB2-expressing cancer cells *in vitro*. In contrast, an antagonistic effect of CI-1033 and an anti-ErbB2 antibody, Herceptin, is disclosed.

[00013] Thus, while there has been a great deal of research interest in the benzoquinone ansamycins, particularly geldanamycin and 17-AAG, there remains a need for effective therapeutic regimens to treat cancer or other diseases or conditions characterized by undesired cellular hyperproliferation using such compounds, whether alone or in combination with other agents. If water-soluble benzoquinone ansamycins were available, such compounds might be more readily formulated and more effective in clinical treatment without dangerous hepatotoxicity. If effective therapeutic treatment regimens were available for administering the benzoquinone ansamycins with other proven anti-cancer compounds, there could be new and more effective means of treating cancer. If the potential of using a benzoquinone ansamycin to lower the effective dose of another anti-cancer agent could be realized, then not only could less expensive therapies be made available (since less drug would need to be administered) but also, and more importantly, one could use drugs that have previously not been useful in chemotherapy due to their narrow therapeutic window. Thus, there is an unmet need for synergists of anti-cancer compounds that allow for administration of doses significantly below the maximum tolerated dose while maintaining therapeutic effectiveness, along with appropriate dosing schedules for combination therapy. The present invention meets such needs in that it provides novel benzoquinone ansamycins and provides methods for using these novel compounds as well as known compounds in single-agent and combination therapies for the treatment of cancer and other diseases or conditions characterized by undesired cellular hyperproliferation.

## SUMMARY OF THE INVENTION

[00014] The present invention provides compounds, methods for their preparation and intermediates thereto, and methods for the use of these compounds

in the treatment of diseases or conditions characterized by undesired cellular proliferation or hyperproliferation.

5 [00015] In one aspect, the invention provides novel benzoquinone ansamycins related to geldanamycin. These analogs are prepared through chemical manipulation and/or genetic engineering. Compounds having improved solubility properties and compounds having conformations optimized to bind Hsp90 are also provided.

10 [00016] In a second aspect, the invention provides genetically engineered forms of the geldanamycin polyketide synthase biosynthetic gene cluster, vectors comprising said gene clusters, host cells comprising said vectors, and methods for the production of geldanamycin analogs using said host cells.

15 [00017] In a third aspect, the invention provides compositions comprising benzoquinone ansamycins for the treatment of diseases or conditions characterized by undesired cellular proliferation or hyperproliferation. In certain embodiments, the disease is cancer.

20 [00018] In a fourth aspect, the invention provides combination therapies comprising the use of a benzoquinone ansamycin and a second agent for use in the treatment of diseases or conditions characterized by undesired cellular hyperproliferation. In certain embodiments, the disease is cancer. In certain embodiments, the second agent is an inhibitor of an Hsp90 client protein. In certain embodiments, the second agent is a protein kinase inhibitor. In certain embodiments, the second agent is a microtubule stabilizing agent. In certain embodiments, the second agent is a cytotoxic drug. In one embodiment, the second agent has been approved by the Federal Drug Administration as a stand-alone agent  
25 for the treatment of cancer. In another embodiment, the second agent has not entered or has entered but not progressed through clinical trials in the United States due to overt toxicity or narrow therapeutic window.

30 [00019] In a fifth aspect, the invention provides methods for preventing undesired cell adhesion and growth on devices for *in vivo* use. These devices include stents, catheters, prostheses and the like.

## BRIEF DESCRIPTION OF THE DRAWINGS

[00020] Figure 1 shows the structures of various naturally-occurring benzoquinone ansamycins as well as 17-AAG and 17-DMAG.

5 [00021] Figure 2 shows particular embodiments of the compounds having formula (I) having groups with solubilizing functionalities.

[00022] Figure 3 shows the results of treating SKBr3 cells with a benzoquinone ansamycin and the protein kinase inhibitor Iressa according to the methods of the present invention. Panel A shows results with 17-AAG. Panel B  
10 shows results with 17-DMAG.

[00023] Figure 4 shows the structures of representative protein kinase inhibitors.

[00024] Figure 5 shows the results of treating H358 cells with 17-AAG and the microtubule stabilizing agent paclitaxel according to the methods of the present  
15 invention.

## DETAILED DESCRIPTION OF THE INVENTION

[00025] The present invention provides compounds, intermediates thereto,  
20 and methods for the use of these compounds in the treatment of diseases or conditions characterized by undesired cellular hyperproliferation.

[00026] Statements regarding the scope of the present invention and definitions of terms used herein are listed below. The definitions apply to the terms as they are used throughout this specification, unless otherwise limited in specific  
25 instances, either individually or as part of a larger group.

[00027] Where stereochemistry is not specifically indicated, all stereoisomers of the inventive compounds are included within the scope of the invention, as pure compounds as well as mixtures thereof. Unless otherwise indicated, individual enantiomers, diastereomers, geometrical isomers, and  
30 combinations and mixtures thereof are all encompassed by the present invention. Polymorphic crystalline forms and solvates are also encompassed within the scope of this invention.



[00028] Protected forms of the inventive compounds are included within the scope of this invention. A variety of protecting groups are disclosed, for example, in T.H. Greene and P.G.M. Wuts, Protective Groups in Organic Synthesis, Third Edition, John Wiley & Sons, New York (1999). The present invention includes within its scope prodrugs of the active compounds of this invention. Such prodrugs are in general functional derivatives of the compounds that are readily convertible *in vivo* into the required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various disorders described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to a subject in need thereof. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs," H. Bundgaard ed., Elsevier, 1985.

[00029] As used herein, the terms "benzoquinone ansamycin" refers to a compound comprising a benzoquinone nucleus connected at two non-adjacent positions by a macrocyclic lactam. Specific examples of naturally-occurring benzoquinone ansamycins include but are not limited to geldanamycin, herbimycin, macbecin, mycotrienes, and ansamitocin. The term "geldanamycin analog" refers to a type of benzoquinone ansamycin that can be derived from geldanamycin by chemical manipulation or by manipulation of the geldanamycin biosynthetic gene cluster, such as 17-allylamino-17-desmethoxygeldanamycin (17-AAG), 17-(2-dimethylaminoethyl)amino-17-desmethoxygeldanamycin (17-DMAG), or a compound having a structure shown in formula (I).

[00030] As used herein, the term "aliphatic" refers to saturated and non-aromatic unsaturated straight chain, branched chain, cyclic, or polycyclic hydrocarbons. Illustrative examples of aliphatic groups include alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, and cycloalkynyl groups. The term "alkyl" refers to a straight or branched chain saturated hydrocarbon substituent. "Alkenyl" refers to a straight or branched chain hydrocarbon substituent with at least one carbon-carbon double bond. "Alkynyl" refers to a straight or branched chain hydrocarbon substituent with at least one carbon-carbon triple bond.

[00031] The term "aryl" refers to monocyclic or polycyclic groups having at least one aromatic ring structure that include preferably one to fourteen carbon atoms. Illustrative examples of aryl groups include but are not limited to: naphthyl, phenyl, tetrahydronaphthyl, and the like.

5 [00032] The term "heteroaryl" refers to monocyclic or polycyclic groups having at least one aromatic ring structure and that include one or more heteroatoms and preferably one to fourteen carbon atoms. Illustrative examples of heteroaryl groups include but are not limited to: furanyl, imidazolyl, indanyl, indolyl, indazolyl, isoxazolyl, isoquinolyl, oxazolyl, oxadiazolyl, pyrazinyl, pyridyl, 10 pyrimidinyl, pyrrolyl, pyrazolyl, quinolyl, quinoxalyl, tetrazolyl, thiazolyl, thienyl, and the like.

[00033] The aliphatic, aryl, and heteroaryl moieties may be substituted with one or more substituents, preferably from one to five substituents, more preferably from one to three substituents, and most preferably from one to two substituents, and 15 as such are referred to as "substituted aliphatic," "substituted aryl," and "substituted heteroaryl." The definition of any substituent or variable at a particular location in a molecule is independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of this invention can be selected by one of ordinary skill in the art to provide compounds 20 that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth herein. Examples of suitable substituents include but are not limited to: aliphatic, haloaliphatic, halogen, aryl, heteroaryl, hydroxy, alkoxy, aryloxy, azido, thio, alkylthio, arylthio, amino, alkylamino, arylamino, acyl, carbamoyl, sulfonamido, nitro, cyano, carboxy, 25 guanidine, and the like.

[00034] The term "haloaliphatic" refers to a substituted aliphatic group substituted by one or more halogens.

[00035] The terms "halo," "halogen," or "halide" refer to fluorine, chlorine, bromine, and iodine.

30 [00036] The term "acyl" refers to  $-C(=O)R$ , where R is an aliphatic group.

[00037] The term "alkoxy" refers to  $-OR$ , where R is an aliphatic group.

[00038] The term "aryloxy" refers to  $-OR$ , where R is an aryl group.

[00039] The term "carbamoyl" refers to  $-O(C=O)NRR'$ , where R and R' are independently H, aliphatic, or aryl groups.

[00040] The term "alkylamino" refers to  $-NHR$ , where R is an alkyl group. The term "dialkylamino" refers to  $-NRR'$ , where both R and R' are alkyl groups.

5 [00041] The term "hydroxyalkyl" refers to  $-R-OH$ , where R is an aliphatic group.

[00042] The term "aminoalkyl" refers to  $-R-NH_2$ , where R is an aliphatic group. The term "alkylaminoalkyl" refers to  $-R-NH-R'$ , where both R and R' are aliphatic groups. The term "dialkylaminoalkyl" refers to  $-R-N(R')-R''$ , where R, R', and R'' are aliphatic groups. The term "arylaminoalkyl" refers to  $-R-NH-R'$ , where R is an aliphatic and R' is an aryl group.

10

[00043] The term "oxo" refers to a carbonyl oxygen ( $=O$ ).

[00044] The term "isolated" as used herein means that the isolated material is in a preparation in which said material forms a major component of the preparation, such as constituting about 50%, about 60%, about 70%, about 80%, about 90%, about 95%, about 99%, or more by weight of the components in the preparation.

15

[00045] The term "subject" as used herein, refers to an animal, typically a mammal or a human, that has been the object of treatment, observation, and/or experiment. When the term is used in conjunction with administration of a compound or drug, then the subject has been the object of treatment, observation, and/or administration of the compound or drug.

20

[00046] The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a cell culture, tissue system, animal, or human that is being sought by a researcher, veterinarian, clinician, or physician, which includes alleviation of the symptoms of the disease, condition, or disorder being treated.

25

[00047] The term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product that results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

30

[00048] The term “pharmaceutically acceptable salt” refers to a salt of one or more compounds. Suitable pharmaceutically acceptable salts of compounds include acid addition salts which may, for example, be formed by mixing a solution of the compound with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, hydrobromic acid, sulfuric acid, fumaric acid, maleic acid, succinic acid, benzoic acid, acetic acid, citric acid, tartaric acid, phosphoric acid, carbonic acid, or the like. Where the compounds carry one or more acidic moieties, pharmaceutically acceptable salts may be formed by treatment of a solution of the compound with a solution of a pharmaceutically acceptable base, such as lithium hydroxide, sodium hydroxide, potassium hydroxide, tetraalkylammonium hydroxide, lithium carbonate, sodium carbonate, potassium carbonate, ammonia, alkylamines, or the like.

[00049] The term “pharmaceutically acceptable carrier” refers to a medium that is used to prepare a desired dosage form of a compound. A pharmaceutically acceptable carrier can include one or more solvents, diluents, or other liquid vehicles; dispersion or suspension aids; surface active agents; isotonic agents; thickening or emulsifying agents; preservatives; solid binders; lubricants; and the like. Remington’s Pharmaceutical Sciences, Fifteenth Edition, E.W. Martin (Mack Publishing Co., Easton, PA, 1975) and Handbook of Pharmaceutical Excipients, Third Edition, A.H. Kibbe ed. (American Pharmaceutical Assoc. 2000), disclose various carriers used in formulating pharmaceutical compositions and known techniques for the preparation thereof.

[00050] The term “pharmaceutically acceptable ester” refers to an ester that hydrolyzes under physiologically relevant conditions to produce a compound or a salt thereof. Illustrative examples of suitable ester groups include but are not limited to formates, acetates, propionates, butyrates, succinates, and ethylsuccinates.

[00051] The term “client protein” refers to a protein that interacts with a chaperone, for example Hsp90. In one aspect, this interaction with a chaperone is useful or required either for proper folding or for stabilization and maintenance. In another aspect, the chaperone forms the core of a functional receptor complex. In both of these aspects, the interaction with the chaperone may be direct or mediated through one or more other proteins. Table 1 below provides an illustrative list of the

client proteins of Hsp90. The term "clientele" refers to the complete set of client proteins for a chaperone.


**Table 1. Illustrative list of Hsp90 client proteins**

5	Client protein	Function
	<i>Steroid Hormone Receptors</i>	
	Glucocorticoid receptor	ligand-mediated gene transcription
	Estrogen receptor	ligand-mediated gene transcription
	Androgen receptor	ligand-mediated gene transcription
10	Progesterone receptor	ligand-mediated gene transcription
	<i>Protein Kinases</i>	
	c-SRC, v-SRC	signal transduction
	LCK	T-cell development & function
	WEE1	Cell cycle regulation (G2)
15	MYT1	Cell cycle regulation (G2)
	ErbB2 (Her-2)	Signal transduction
	EGFR (ErbB1)	Signal transduction
	FPS/FES	Cell proliferation
	c-RAF-1, v-RAF-1	MAPK signaling
20	MEK	MAPK signaling
	Casein kinase 2	pleiotropic kinase
	CDK4	Cell cycle regulation (G1)
	AKT (PKB)	PI3 kinase signaling
	Death domain kinase RIP	TNF-mediated necrosis
25	Bcr-ABL	myeloid leukemia pathogenesis
	PIM-1	cytokine-mediated proliferation
	MOK	MAPK signaling
	Polo-1 kinase (PLK)	Cell cycle regulation (G2/M)
	Focal adhesion kinase (FAK)	actin-based cell motility
30	c-MET	HGF/SF-MET motility signaling
	eIF2 kinase	transcriptional regulation
	<i>Other Client Proteins</i>	
	Mutant p53	cell cycle checkpoint protein mutant
35	Hepatitis B reverse transcriptase	viral transcription
	hTERT (telomerase subunit)	cell mortality, senescence
	$\beta\gamma$ subunits of trimeric G proteins	signal transduction
	endothelial NOS	NO synthesis
	calcineurin	$\text{Ca}^{2+}$ -dependent signaling
40	tubulin	microtubule formation
	HIF-1 $\alpha$	hypoxia-induced angiogenesis
	Retinoblastoma protein	cell cycle regulation (G1/S)
	Tumor necrosis factor receptor 1	TNF-mediated apoptosis
	Cystic fibrosis transmembrane	
45	conductance regulator	Ion channel/cystic fibrosis
	Immunoglobulin chains	Immune response

Fanconi anemia protein	hematopoiesis
Apoprotein B	atherosclerosis
Aryl hydrocarbon receptor	gene transcription
SV40 T antigen	viral oncogene

5

[00052] In one aspect of the invention, geldanamycin analogs having the  
 10 formula (I) are provided:



I

wherein  $R^1$  is MeO,  $(CH_2)_3N$  or  $R^9-NH$ , wherein  $R^9$  is selected from the group consisting of H, substituted or unsubstituted  $C_1-C_6$  alkyl, substituted or unsubstituted  $C_1-C_6$  alkenyl, substituted or unsubstituted  $C_1-C_6$  alkynyl, substituted or unsubstituted  $C_3-C_6$  cycloalkyl, piperidinyl, N-alkylpiperidinyl, hexahydropyranyl, furfuryl, tetrahydrofurfuryl, pyrrolidinyl, N-alkylpyrrolidinyl, piperazinylamino, N-alkylpiperazinyl, morpholinyl, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-methyl-pyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl, or  $R^6$  is H and  $R^1$  and  $R^5$  taken together form a group of the formula  $NH-Z-O$ , wherein Z is a linker comprised of from 1 to 6 carbon atoms and 0 to 2 nitrogen atoms and wherein the O is attached at the position of  $R^5$ ;  $R^2$  is selected from the group consisting of H, halogen,  $OR^{10}$ ,  $NHR^{10}$ ,  $SR^{10}$ , aryl, and heteroaryl, wherein  $R^{10}$  is selected from the group consisting of substituted or unsubstituted  $C_1-C_6$  alkyl, substituted or unsubstituted  $C_1-C_6$  alkenyl, substituted or unsubstituted  $C_1-C_6$  alkynyl, and substituted or unsubstituted  $C_3-C_6$  cycloalkyl;  $R^3$  is H, OH, or OMe;  $R^4$  is H or Me;  $R^5$  is OH or  $O-C(=O)-CH_2NH_2$  and  $R^6$  is H, or  $R^5$  and  $R^6$  taken together

form =O or =N-OR<sup>11</sup>, wherein R<sup>11</sup> is selected from the group consisting of H, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkenyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkynyl, substituted or unsubstituted C<sub>3</sub>-C<sub>6</sub> cycloalkyl, aryl, or heteroaryl; R<sup>7</sup> is H and R<sup>8</sup> is H or OH, or R<sup>7</sup> and R<sup>8</sup> taken together form a bond; and X is O or a bond, with the provisos that when R<sup>3</sup> is H, R<sup>4</sup> is Me, and R<sup>7</sup> is H and R<sup>8</sup> is H or R<sup>7</sup> and R<sup>8</sup> taken together form a bond that either R<sup>6</sup> is H and R<sup>1</sup> and R<sup>5</sup> taken together form a group of the formula NH-Z-O, wherein Z is a linker comprised of from 1 to 6 carbon atoms and 0 to 2 nitrogen atoms and wherein the O is attached at the position of R<sup>5</sup>, or that R<sup>1</sup> is (CH<sub>2</sub>)<sub>3</sub>N or R<sup>9</sup>-NH, wherein R<sup>9</sup> is selected from the group consisting of piperidiny, N-alkylpiperidiny, hexahydropyrany, furfuryl, tetrahydrofurfuryl, pyrrolidiny, N-alkylpyrrolidiny, piperazinylamino, N-alkylpiperazinyl, morpholinyl, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-methyl-pyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl; and that when R<sup>3</sup> is H and R<sup>4</sup> is Me that R<sup>7</sup> is H and R<sup>8</sup> is OH.

[00053] In one embodiment, compounds having formula (I) are provided wherein R<sup>1</sup> is (CH<sub>2</sub>)<sub>3</sub>N or R<sup>9</sup>-NH, wherein R<sup>9</sup> is selected from the group consisting of piperidiny, N-alkylpiperidiny, hexahydropyrany, furfuryl, tetrahydrofurfuryl, pyrrolidiny, N-alkylpyrrolidiny, piperazinylamino, N-alkylpiperazinyl, morpholinyl, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-methyl-pyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl; R<sup>2</sup> is selected from the group consisting of H, halogen, OR<sup>10</sup>, NHR<sup>10</sup>, SR<sup>10</sup>, aryl, and heteroaryl, wherein R<sup>10</sup> is selected from the group consisting of substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkenyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkynyl, and substituted or unsubstituted C<sub>3</sub>-C<sub>6</sub> cycloalkyl; R<sup>3</sup> is H, OH, or OMe; R<sup>4</sup> is H or Me; R<sup>5</sup> is OH or O-C(=O)-CH<sub>2</sub>NH<sub>2</sub> and R<sup>6</sup> is H, or R<sup>5</sup> and R<sup>6</sup> taken together form =O or =N-OR<sup>11</sup>, wherein R<sup>11</sup> is selected from the group consisting of H, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkenyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkynyl, substituted or unsubstituted C<sub>3</sub>-C<sub>6</sub> cycloalkyl, aryl, or heteroaryl; R<sup>7</sup> is H and R<sup>8</sup> is H or OH, or R<sup>7</sup> and R<sup>8</sup> taken together form a bond; and X is O or a bond.


[00054] In one embodiment, compounds having formula (I) are provided wherein  $R^1$  is  $(CH_2)_3N$  or  $R^9-NH$ , wherein  $R^9$  is selected from the group consisting of piperidinyl, N-alkylpiperidinyl, hexahydropyranyl, furfuryl, tetrahydrofurfuryl, pyrrolidinyl, N-alkylpyrrolidinyl, piperazinylamino, N-alkylpiperazinyl, morpholinyl, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-methyl-pyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl;  $R^2$  is H;  $R^3$  is H, OH, or OMe;  $R^4$  is H or Me;  $R^5$  is OH or  $O-C(=O)-CH_2NH_2$  and  $R^6$  is H, or  $R^5$  and  $R^6$  taken together form  $=O$  or  $=N-OR^{11}$ , wherein  $R^{11}$  is selected from the group consisting of H, substituted or unsubstituted  $C_1-C_6$  alkyl, substituted or unsubstituted  $C_1-C_6$  alkenyl, substituted or unsubstituted  $C_1-C_6$  alkynyl, substituted or unsubstituted  $C_3-C_6$  cycloalkyl, aryl, or heteroaryl;  $R^7$  is H and  $R^8$  is H or OH, or  $R^7$  and  $R^8$  taken together form a bond; and X is O or a bond.

[00055] In one embodiment, geldanamycin analogs having improved solubility are provided resulting from chemical manipulation of geldanamycin to provide compounds having formula (I) wherein:  $R^1$  is  $(CH_2)_3N$  or  $R^9-NH$ , wherein  $R^9$  is selected from the group consisting of H, substituted or unsubstituted  $C_1-C_6$  alkyl, substituted or unsubstituted  $C_1-C_6$  alkenyl, substituted or unsubstituted  $C_1-C_6$  alkynyl, substituted or unsubstituted  $C_3-C_6$  cycloalkyl, piperidinyl, N-alkylpiperidinyl, hexahydropyranyl, furfuryl, tetrahydrofurfuryl, pyrrolidinyl, N-alkylpyrrolidinyl, piperazinylamino, N-alkylpiperazinyl, morpholinyl, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-methyl-pyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl;  $R^2$  is selected from the group consisting of H, halogen,  $OR^{10}$ ,  $NHR^{10}$ ,  $SR^{10}$ , aryl, and heteroaryl, wherein  $R^{10}$  is selected from the group consisting of substituted or unsubstituted  $C_1-C_6$  alkyl, substituted or unsubstituted  $C_1-C_6$  alkenyl, substituted or unsubstituted  $C_1-C_6$  alkynyl, and substituted or unsubstituted  $C_3-C_6$  cycloalkyl;  $R^3$  is H;  $R^4$  is methyl;  $R^5$  is OH or  $O-C(=O)-CH_2NH_2$  and  $R^6$  is H, or  $R^5$  and  $R^6$  taken together form  $=O$  or  $=N-OH$ ;  $R^7$  and  $R^8$  are H, or  $R^7$  and  $R^8$  taken together form a bond; and X is O or a bond.



[00056] In another embodiment of the invention, compounds having formula (I) are provided wherein:  $R^1$  is  $R^9$ -NH, wherein  $R^9$  is selected from the group consisting of ethyl, 2-(dimethylamino)ethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2-(N-methylpyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, 3-(4-morpholino)-1-propyl, 3-(dimethylamino)-1-propyl, 3-(dimethylamino)-2-propyl, 2-(dimethylamino)-1-propyl, and cyclopropyl-methyl;  $R^2$  is H;  $R^3$  is H;  $R^4$  is methyl;  $R^5$  is OH or  $O-C(=O)-CH_2NH_2$  and  $R^6$  is H, or  $R^5$  and  $R^6$  taken together form  $=O$  or  $=N-OH$ ;  $R^7$  and  $R^8$  are H, or  $R^7$  and  $R^8$  taken together form a bond; and X is O or a bond.


[00057] In another embodiment of the invention, compounds having the structures:



are provided.

[00058] In another embodiment, geldanamycin analogs having formula (I) are provided wherein  $R^1$  is OMe;  $R^2$  is H;  $R^3$  is H, OH, or OMe;  $R^4$  is H or methyl;  $R^5$  is OH and  $R^6$  is H;  $R^7$  is H and  $R^8$  is H or OH, or  $R^7$  and  $R^8$  taken together form a bond; and X is a bond, with the proviso that geldanamycin and 4,5-dihydrogeldanamycin are not included.


[00059] Other embodiments of the invention provide compounds having the formulas:



[00060] In other embodiments of the invention, the geldanamycin analogs described above serve as starting materials for chemical addition of solubilizing groups. In one embodiment, 15-hydroxygeldanamycin is derivatized to provide compounds having formula (I) wherein:  $R^1$  is  $(CH_2)_3N$  or  $R^9-NH$ , wherein  $R^9$  is selected from the group consisting of H, substituted or unsubstituted  $C_1-C_6$  alkyl, substituted or unsubstituted  $C_1-C_6$  alkenyl, substituted or unsubstituted  $C_1-C_6$  alkynyl, substituted or unsubstituted  $C_3-C_6$  cycloalkyl, piperidiny, N-alkylpiperidiny, hexahydropyrany, furfuryl, tetrahydro-furfuryl, pyrrolidiny, N-alkylpyrrolidiny, piperazinylamino, N-alkylpiperazinyl, morpholiny, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-methyl-pyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl;  $R^3$  is OH;  $R^4$  is methyl;  $R^5$  is OH or  $O-C(=O)-CH_2NH_2$  and  $R^6$  is H, or  $R^5$  and  $R^6$  taken together form  $=O$  or  $=N-OH$ ; and  $R^7$  and  $R^8$  taken together form a bond; and X is O or a bond.

[00061] In another embodiment of the invention, 15-hydroxygeldanamycin is derivatized to provide compounds having formula (I) wherein:  $R^1$  is  $(CH_2)_3N$  or  $R^9-NH$ , wherein  $R^9$  is selected from the group consisting of allyl, ethyl, 2-(dimethylamino)ethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2-(N-methylpyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, 3-(4-morpholino)-1-propyl, 3-(dimethylamino)-1-propyl, 3-(dimethylamino)-2-propyl, 2-(dimethylamino)-1-propyl, and cyclopropylmethyl;  $R^2$  is H;  $R^3$  is OH;  $R^4$  is methyl;  $R^5$  is OH or  $O-C(=O)-CH_2NH_2$  and  $R^6$  is H, or  $R^5$  and  $R^6$  taken together form  $=O$  or  $=N-OH$ ;  $R^7$  and  $R^8$  taken together form a bond; and X is O or a bond.

[00062] In another embodiment, 15-hydroxygeldanamycin analogs having the formulas




are provided.

[00063] In one embodiment, 28-desmethylgeldanamycin is derivatized to provide compounds having formula (I) wherein:  $R^1$  is  $(CH_2)_3N$  or  $R^9-NH$ , wherein  $R^9$  is selected from the group consisting of H, substituted or unsubstituted  $C_1-C_6$  alkyl, substituted or unsubstituted  $C_1-C_6$  alkenyl, substituted or unsubstituted  $C_1-C_6$  alkynyl, substituted or unsubstituted  $C_3-C_6$  cycloalkyl, piperidinyl, N-alkylpiperidinyl, hexahydropyranyl, furfuryl, tetrahydro-furfuryl, pyrrolidinyl, N-alkylpyrrolidinyl, piperazinylamino, N-alkylpiperazinyl, morpholinyl, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-methylpyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl;  $R^2$  is selected from the group consisting of H, halogen,  $OR^{10}$ ,  $NHR^{10}$ ,  $SR^{10}$ , aryl, and heteroaryl, wherein  $R^{10}$  is selected from the group consisting of substituted or unsubstituted  $C_1-$

C<sub>6</sub> alkyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkenyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkynyl, and substituted or unsubstituted C<sub>3</sub>-C<sub>6</sub> cycloalkyl; R<sup>3</sup> is H; R<sup>4</sup> is H; R<sup>5</sup> is OH or O-C(=O)-CH<sub>2</sub>NH<sub>2</sub> and R<sup>6</sup> is H, or R<sup>5</sup> and R<sup>6</sup> taken together form =O or =N-OH; and R<sup>7</sup> and R<sup>8</sup> taken together form a bond; and X is O or a bond.

5           **[00064]** In another embodiment of the invention, 28-desmethylgeldanamycin is derivatized to provide compounds having formula (I) wherein: R<sup>1</sup> is (CH<sub>2</sub>)<sub>3</sub>N or R<sup>9</sup>-NH, wherein R<sup>9</sup> is selected from the group consisting of allyl, ethyl, 2-(dimethylamino)ethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2-(N-methylpyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl,  
10       2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, 3-(4-morpholino)-1-propyl, 3-(dimethylamino)-1-propyl, 3-(dimethylamino)-2-propyl, 2-(dimethylamino)-1-propyl, and cyclopropylmethyl; R<sup>2</sup> is H; R<sup>3</sup> is H; R<sup>4</sup> is H; R<sup>5</sup> is OH or O-C(=O)-CH<sub>2</sub>NH<sub>2</sub> and R<sup>6</sup> is H, or R<sup>5</sup> and R<sup>6</sup> taken together form =O or =N-OH; R<sup>7</sup> and R<sup>8</sup> taken together form a bond; and X is O or a bond.

15           **[00065]** In another embodiment, 28-desmethylgeldanamycin analogs having the formulas




are provided.

20           **[00066]** In another embodiment, 4,5-dihydro-5-hydroxygeldanamycin is derivatized to provide compounds having formula (I) wherein: R<sup>1</sup> is (CH<sub>2</sub>)<sub>3</sub>N or R<sup>9</sup>-NH, wherein R<sup>9</sup> is selected from the group consisting of H, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkenyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkynyl, substituted or unsubstituted C<sub>3</sub>-C<sub>6</sub> cycloalkyl, piperidinyl, N-alkylpiperidinyl, hexahydropyranyl, furfuryl, tetrahydro-furfuryl,  
25       pyrrolidinyl, N-alkylpyrrolidinyl, piperazinylamino, N-alkylpiperazinyl, morpholinyl, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-

methylpyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl;  $R^2$  is selected from the group consisting of H, halogen,  $OR^{10}$ ,  $NHR^{10}$ ,  $SR^{10}$ , aryl, and heteroaryl, wherein  $R^{10}$  is selected from the group consisting of substituted or unsubstituted  $C_1$ - $C_6$  alkyl, substituted or unsubstituted  $C_1$ - $C_6$  alkenyl, substituted or unsubstituted  $C_1$ - $C_6$  alkynyl, and substituted or unsubstituted  $C_3$ - $C_6$  cycloalkyl;  $R^3$  is H;  $R^4$  is methyl;  $R^5$  is OH or  $O-C(=O)-CH_2NH_2$  and  $R^6$  is H, or  $R^5$  and  $R^6$  taken together form  $=O$  or  $=N-OH$ ; and  $R^7$  is H;  $R^8$  is OH; and X is O or a bond.

[00067] In another embodiment of the invention, 4,5-dihydro-5-hydroxygeldanamycin is derivatized to provide compounds having formula (I) wherein:  $R^1$  is  $(CH_2)_3N$  or  $R^9-NH$ , wherein  $R^9$  is selected from the group consisting of allyl, ethyl, 2-(dimethylamino)ethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2-(N-methylpyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, 3-(4-morpholino)-1-propyl, 3-(dimethylamino)-1-propyl, 3-(dimethylamino)-2-propyl, 2-(dimethylamino)-1-propyl, and cyclopropylmethyl;  $R^2$  is H;  $R^3$  is H;  $R^4$  is methyl;  $R^5$  is OH or  $O-C(=O)-CH_2NH_2$  and  $R^6$  is H, or  $R^5$  and  $R^6$  taken together form  $=O$  or  $=N-OH$ ;  $R^7$  is H;  $R^8$  is OH; and X is O or a bond.

[00068] In another embodiment, 4,5-dihydro-5-hydroxygeldanamycin analogs having the formulas



are provided.


[00069] In another aspect of the invention, conformationally constrained geldanamycin analogs are provided. In one embodiment, compounds having the formula (I) are provided wherein  $R^1$  and  $R^5$  taken together form a group of the formula  $NH-Z-O$ , wherein Z is a linker comprised of from 1 to 6 carbon atoms and 0

to 2 nitrogen atoms and wherein the O is attached at the position of R<sup>5</sup>; R<sup>2</sup> is selected from the group consisting of H, halogen, OR<sup>10</sup>, NHR<sup>10</sup>, SR<sup>10</sup>, aryl, and heteroaryl, wherein R<sup>10</sup> is selected from the group consisting of substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkenyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkynyl, and substituted or unsubstituted C<sub>3</sub>-C<sub>6</sub> cycloalkyl; R<sup>3</sup> is H or OH; R<sup>4</sup> is H or methyl; R<sup>7</sup> is H and R<sup>8</sup> is H or OH, or R<sup>7</sup> and R<sup>8</sup> taken together form a bond; and X is O or a bond.

[00070] In one embodiment of the invention, compounds having formula (I) are provided wherein R<sup>1</sup> and R<sup>5</sup> taken together form a group of the formula NH-(CH<sub>2</sub>)<sub>4</sub>-O, NH-CH<sub>2</sub>CH=CHCH<sub>2</sub>-O, or NH-CH<sub>2</sub>CCCH<sub>2</sub>-O; R<sup>2</sup> is H; R<sup>3</sup> is H, OH, or OMe; R<sup>4</sup> is H or methyl; R<sup>7</sup> is H and R<sup>8</sup> is H or OH, or R<sup>7</sup> and R<sup>8</sup> taken together form a bond; and X is O or a bond.

[00071] In another embodiment of the invention, compounds having formula (I) are provided wherein R<sup>1</sup> and R<sup>5</sup> taken together form a group of the formula NH-CH<sub>2</sub>CH=CHCH<sub>2</sub>-O; R<sup>2</sup> is H; R<sup>3</sup> is H, OH, or OMe; R<sup>4</sup> is H or methyl; R<sup>7</sup> is H and R<sup>8</sup> is H or OH, or R<sup>7</sup> and R<sup>8</sup> taken together form a bond; and X is O or a bond.


[00072] In another embodiment of the invention, compounds having formula (I) are provided having the structures:



[00073] In another embodiment, the water-soluble analogs described above are subjected to conformational constraint to provide geldanamycin analogs having both improved solubility and improved specificity for Hsp90. Poor water solubility is a major factor limiting the clinical usefulness of geldanamycin and 17-AAG. Improvements in water solubility of a compound can be achieved according to the methods of the present invention either by addition of groups containing solubilizing functionalities to the compound or by removal of hydrophobic groups from the compound, so as to decrease the lipophilicity of the compound. Typical groups containing solubilizing functionalities are shown in Figure 2 and include but are not limited to: 2-(dimethylaminoethyl)amino, piperidinyl, N-alkylpiperidinyl, hexahydropyranyl, furfuryl, tetrahydrofurfuryl, pyrrolidinyl, N-alkylpyrrolidinyl, piperazinylamino, N-alkylpiperazinyl, morpholinyl, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-methylpyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl.

[00074] Solubilizing groups are added to the geldanamycin analog by reaction of geldanamycin with amines, which results in the displacement of the 17-methoxy group by the amine as illustrated in Scheme 1 and exemplified in Example 1 (Schnur *et al.* (1995) "Inhibition of the oncogene product p185<sup>erbB-2</sup> in Vitro and in Vivo by Geldanamycin and Dihydrogeldanamycin Derivatives," *J. Med. Chem.* 38, 3806-3812; Schnur *et al.* (1995) "*erbB-2* Oncogene Inhibition by Geldanamycin Derivatives: Synthesis, mechanism of Action, and Structure-Activity relationships," *J. Med. Chem.* 38, 3813-3820; Schnur *et al.*, "Ansamycin derivatives as antioncogene and anticancer agents," U.S. Patent 5,932,655; all of which are incorporated herein by reference). Typical amines containing solubilizing functionalities include 2-(dimethylamino)-ethylamine, 4-aminopiperidine, 4-amino-1-methylpiperidine, 4-aminohexahydropyran, furfurylamine, tetrahydrofurfurylamine, 3-(aminomethyl)-tetrahydrofuran, 2-(aminomethyl)pyrrolidine, 2-(aminomethyl)-1-methylpyrrolidine, 1-methylpiperazine, morpholine, 1-methyl-2-(aminomethyl)aziridine, 1-(2-aminoethyl)-1-azabicyclo-[1.3.0]hexane, 1-(2-aminoethyl)piperazine, 4-(2-aminoethyl)morpholine, 1-(2-amino-ethyl)pyrrolidine, 2-(2-aminoethyl)pyridine, 2-fluoroethylamine, 2,2-difluoroethylamine, and the like.

SCHEME 1




[00075] Similar solubilizing groups can be introduced by treatment of 19-bromo-geldanamycin or analogs with an amine containing a solubilizing substituent in accordance with the methods of the present invention, resulting in a 19-amino-substituted geldanamycin analog. The 19-bromo derivative is formed upon treatment of the geldanamycin analog with a suitable brominating reagent, such as pyridinium bromide perbromide (Schnur *et al.* 1995, *J. Med. Chem.* 38, 3806-3812;




incorporated herein by reference). Reaction of 19-bromogeldanamycin with an arylboronic acid in the presence of a palladium catalyst according to the method of the present invention gives 19-aryl substituted geldanamycins as illustrated in Scheme 2. Boronic acids such as phenylboronic acid, (4-dimethyl-amino)phenylboronic acid, 4-pyridylboronic acid, 4-methoxyphenylboronic acid, 2-furylboronic acid, and 4-(dimethylaminomethyl)-phenylboronic acid and the like can also be used.

SCHEME 2



[00076] In another embodiment of the invention, the geldanamycin analogs are oxidized to produce the corresponding 11-oxogeldanamycin analogs as illustrated in Scheme 3.

SCHEME 3



The 11-oxogeldanamycin analog resulting from oxidation of the 11-OH are converted into the 11-oximino analogs by reaction with hydroxylamine or an alkyoxylamine.

[00077] In another embodiment of the invention, the geldanamycin analog is treated with a peroxyacid, for example 3-chloroperoxybenzoic acid (mCPBA), to produce the 8,9-epoxide, as illustrated in Scheme 3.

In another aspect, the invention provides genetically engineered forms of the geldanamycin polyketide synthase biosynthetic gene cluster, vectors comprising said gene clusters, host cells comprising said vectors, and methods for the production of geldanamycin analogs using said host cells.

[00078] In one embodiment of the invention, substitution of the acyltransferase domain in module 1 of the geldanamycin PKS gene with one specific for malonyl-CoA instead of 2-methylmalonyl-CoA results in formation of 28-desmethyl-geldanamycin. The domain swap is created by introducing a malonyl-CoA specific acyltransferase domain from a heterologous PKS gene, for example from the rapamycin, tylosin, or FK520 PKS genes or the like, into the geldanamycin

PKS locus by homologous recombination into a strain which produces geldanamycin, aided by a selectable antibiotic resistance gene, then isolating the recombinants resulting from double crossover events in which the wild-type acyltransferase domain is replaced with one specific for malonyl-CoA. Details of this are provided below in Example 5.

[00079] In another embodiment of the invention, the acyltransferase domain in module 1 of the geldanamycin PKS gene is mutagenized according to the methods described in Reeves *et al.*, "Alteration of the substrate specificity of a modular polyketide synthase acyltransferase domain through site-directed mutagenesis," *Biochemistry* 2001, 40: 15464-15470, and in U.S. patent application serial no. 60/310,730, entitled "Alteration of the substrate specificity of a modular PKS AT domain," which is incorporated herein by reference. Details of this are provided below in Example 6.

[00080] In another embodiment of the invention, the coding sequence for the reduction cassette of module 6, which has both DH and KR domains, is replaced with a coding sequence for a reduction cassette that has only a KR domain. Details of this are provided below in Example 7.

[00081] In another embodiment of the invention, inactivation of the dehydraase domain in module 6 of the geldanamycin PKS gene by site-specific mutation of the wild-type domain in accord with the methods of the present invention results in production of 4,5-dihydro-5-hydroxygeldanamycin. Details of this are provided below in Example 8.

[00082] In another embodiment of the invention, a substantial portion of the nucleotide sequence in module 6 between the end of the AT domain is deleted to provide 4,5-dihydro-5-hydroxy-geldanamycin. Details of this are provided below in Example 9.

[00083] In another embodiment of the invention, the dehydratase domain of module 1 is replaced or inactivated as described above for module 6 to provide 15-hydroxy-geldanamycin. Details of this are provided below in Example 10.

[00084] In another embodiment of the invention, inactivation of the dehydratase domain in module 1 of the geldanamycin PKS gene by site-specific mutation of the wild-type domain in accord with the methods of the present

invention results in production of 15-hydroxygeldanamycin. Details of this are provided below in Example 11.

5           [00085]   It is also possible to express the geldanamycin gene cluster or mutated versions of the geldanamycin gene cluster prepared according to the methods of the invention in host cells other than the native geldanamycin producer. Methods for heterologous expression of PKS genes and host cells suitable for expression of these genes and production of polyketides are described, for example, in U.S. Patent Nos. 5,843,718 and 5,830,750; PCT publications WO 01/31035 and WO 01/27306; and U.S. patent application serial nos. 10/087,451; 60/\_\_\_\_,\_\_\_\_, 10           entitled "Process and Metabolic Strategies for Improved Production of *E. coli* derived 6-deoxyerythronolide B," by inventors Pfeifer and Khosla (atty docket no. 30226.00); and 60/\_\_\_\_,\_\_\_\_, entitled "Metabolic Pathways for Starter Units in Polyketide Biosynthesis in *E. Coli*" by inventors Kealey, Dayem, and Santi (atty docket no. 30088.00); each of which is incorporated herein by reference.

15           [00086]   Inactivation of dehydratase domains in accord with the methods of the present invention may also be obtained through random mutagenesis of the organism that normally produces geldanamycin. In this instance, spores of the producing organism can be either treated with a chemical mutagen, for example 1-methyl-3-nitro-1-nitrosoguanidine (MNNG), dimethylsulfate, or the like, or with 20           mutagenic levels of radiation, for example ultraviolet radiation. The surviving spores are then allowed to grow on a suitable medium, and the resulting cultures are analyzed, for example by LC-mass spectrometry, for the presence of the desired new geldanamycin analog. Methods for the random mutagenesis of *Streptomyces* are described in Kieser *et al*, "Practical Streptomyces Genetics," The John Innes 25           Foundation, Norwich (2000), which is incorporated herein by reference.


          [00087]   In accordance with the methods of the present invention, replacement of other acyltransferase domains in the geldanamycin PKS can be used to generate the respective desmethyl or desmethoxy analogs, and replacement of other dehydratase domains with inactive versions can be used to generate the 30           corresponding dihydro-hydroxy analogs. Such analogs are expected to be more water-soluble, as they have fewer lipophilic substituents (28-

desmethylgeldanamycin) or have additional hydrophilic substituents (4,5-dihydro-5-hydroxygeldanamycin or 15-hydroxygeldanamycin).

[00088] By using the geldanamycin analogs produced by genetic engineering of the geldanamycin PKS in accord with the methods of the present invention as described above, the afore-mentioned chemical transformations can be used to convert the analogs into more water-soluble, more potent, more specific inhibitors of Hsp90. Such compounds thus may overcome the necessity of using toxic vehicles such as Cremophore® in their administration, and show improved selectivity and reduced side-effects. In one embodiment of the invention, a geldanamycin analog prepared by genetic engineering is reacted with an amine as illustrated in Scheme 1 above.

[00089] Ring-forming olefin metathesis of 11-O-allyl-17-allylamino-17-desmethoxy-geldanamycin or similar analogs generates conformationally constrained benzoquinone ansamycins structure in accordance with the methods of the present invention. In one embodiment of the invention, treatment of 17-AAG with an allylating reagent, such as allyl *tert*-butyl carbonate and a palladium catalyst, generates 11-O-allyl-17-allylamino-17-desmethoxy-geldanamycin, as illustrated in Scheme 4 and exemplified in Example 2.


SCHEME 4



[00090] Reaction of related geldanamycin analogs resulting from displacement of the 17-methoxy group with amines other than allylamine, e.g., but-3-en-1-ylamine and pent-4-en-1-ylamine, results in formation of the corresponding 17-butenylamino-11-O-allyl and 17-pentenylamino-11-O-allyl analogs in accordance with the methods of the present invention. Treatment of these 11-O-allyl compounds with an olefin metathesis catalyst, such as benzyldiene

bis(tricyclohexylphosphine)ruthenium dichloride, as illustrated in Scheme 5, results in linkage of the 11- and 17-positions through a carbon chain and constraint of the geldanamycin conformation. This is exemplified below in Example 3.


SCHEME 5



[00091] Alteration of the 11-O- and 17-N- groups in accord with the methods of the present invention allows for variation of the linker chain length. Thus, use of 11-O-allyl-17-allylamino-17-desmethoxygeldanamycin results in a 4-carbon linker, use of 11-O-allyl-17-(but-3-en-1-ylamino)-17-desmethoxygeldanamycin results in a 5-carbon linker, and use of 11-O-allyl-17-(pent-4-en-1-ylamino)-17-desmethoxygeldanamycin results in a 6-carbon linker. The carbon-carbon double bond of the linker can optionally be reduced by reduction, e.g. using diimide, to provide a saturated linker.

[00092] An alkynyl linker can be prepared according to the methods of the invention by treating geldanamycin or a geldanamycin analog with a bifunctional alkyne comprising an amino function at one end of the linker and a displaceable function, for example a halogen or sulfonate ester, at the other end. Reaction of geldanamycin or an analog with this bifunctional linker results first in displacement of the 17-methoxy group by the amine. Subsequent base treatment results in alkylation of the 11-hydroxyl as illustrated in Scheme 6.

SCHEME 6



[00093] As described above, alteration of the linker lengths can be achieved through variation in the number of carbon atoms in the chain.

[00094] In another aspect of the invention, a composition comprising a benzoquinone ansamycin is used to treat a disease or condition characterized by undesired cellular proliferation or hyperproliferation. In one embodiment, the disease is cancer. In another embodiment, the disease is stenosis or restenosis. In another embodiment, the disease is psoriasis. In another embodiment, the disease is a neurodegenerative disease. In preferred embodiments, the benzoquinone ansamycin is a compound having formula (I), 17-AAG, or 17-DMAG.

[00095] In another aspect of the invention, a benzoquinone ansamycin is used in combination therapy with a second agent. In one embodiment, the second agent is an inhibitor of an Hsp90 client protein. Suitable Hsp90 client proteins include but are not limited to those listed in Table 1.

[00096] In one embodiment of the invention, a benzoquinone ansamycin is used in combination therapy with a protein kinase inhibitor. Suitable protein kinase inhibitors include but are not limited to the compounds listed in Table 2.

Table 2. Illustrative list of protein kinase inhibitors

	Compound	target	most advanced indication
	<i>Quinazolines and related heterocycles:</i>		
	Iressa (ZD 1839)	EGFR	non-small cell lung cancer
	Tarceva (OSI-774)	EGFR	ovarian cancer
5	GW2016	EGFR	cancer
	CI 1033	EGFR	cancer
	AZD6474	VEGFR	solid tumors
	<i>Phenylamino-pyrimidines:</i>		
	Gleevec (STI-571)	bcr-abl, others	chronic myelogenous leukemia
10	<i>Pyrazolopyrimidines and pyrrolopyrimidines:</i>		
	BIBX 1382	EGFR	cancer
	PKI 166	EGFR	cancer
	<i>Indoles and oxindoles:</i>		
	Semaxanib (SU5416)	flk-1/KDR kinase	advanced colorectal cancer
15	SU5402	FGFR	cancer
	<i>Benzylidene malononitriles (tyrphostins):</i>		
	Tyrphostin 25	EGFR	cancer
	SU101	PDGFR	solid tumors
	Lefunamide (SU0020)	PDGFR	solid tumors
20	<i>Flavones:</i>		
	Flavopiridol	cdk	cancer
	B43-genistein	LYN	leukemia
	<i>Staurosporines:</i>		
	CEP-701	flt-3	prostate, pancreatic cancers
25	CEP-2563	trk	prostate, other cancers
	UCN-01 (NSC638850)		cancer
	LY-333531	PKC $\beta$	diabetic complications
	<i>Antibodies, ribozymes:</i>		
	Herceptin	Her-2	breast cancer
30	Avastin antibody	VEGFR	advanced solid tumors
	2c4 antibody	Her-2	solid tumors
	IMC-1C11	VEGFR	metastatic colorectal cancer
	Cetuximab (C255)	EGFR	head & neck squamous cell cancer
35	ABX-EGF	EGFR	cancer
	TheraCIM (H-R3)	EGFR	metastatic squamous cell carcinoma
	SMART anti-VEGF	VEGFR	relapsed, refractory solid tumors
40	Angiozyme	VEGFR	cancer

[00097] The protein kinase inhibitors listed in Table 2 may be classified according to their chemotypes, including: quinazolines, particularly 4-anilinoquinazolines such as Iressa (AstraZeneca; N-(3-chloro-4-fluorophenyl)-7-methoxy-6-[3-(4-morpholinyl)propoxy]-4-quinazolinamine) and Tarceva (Roche/Genentech; N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-



Quinazolinamine monohydrochloride); phenylamino-pyrimidines such as Gleevec (Novartis; 4-[(4-methyl-1-piperazinyl)methyl]-N-[4-methyl-3-[[4-(3-pyridinyl)-2-pyrimidinyl]amino]phenyl]benzamide); pyrrolo- and pyrazolopyrimidines such as BIBX 1382 (Boehringer Ingelheim; N8-(3-chloro-4-fluorophenyl)-N2-(1-methyl-4-piperidinyl)-pyrimido[5,4-d]pyrimidine-2,8-diamine); indoles and oxindoles such as Semaxinib (Pharmacia; 3-[(3,5-dimethyl-1H-pyrrol-2-yl)methylene]-1,3-dihydro-2H-Indol-2-one); benzylidene malononitriles; flavones such as flavopiridol (Aventis; 2-(2-chlorophenyl)-5,7-dihydroxy-8-[(3S,4R)-3-hydroxy-1-methyl-4-piperidinyl]-4H-1-benzopyran-4-one); staurosporines such as CEP-701 (Cephalon); antibodies such as Herceptin (Genentech); and ribozymes such as Angiozyme (Ribozyme Pharmaceuticals). Structures of representative protein kinase inhibitors are given in Figure 4.

[00098] In another embodiment of the invention, a benzoquinone ansamycin is used in combination therapy with a microtubule stabilizing agent, including paclitaxel, epothilone, discodermolide, and laulimalide. In preferred embodiments, the benzoquinone ansamycin is a compound having the formula (I), 17-AAG, or 17-DMAG.

[00099] In another aspect, the present invention provides combination therapy methods for the treatment of diseases or conditions characterized by undesired cellular proliferation or hyperproliferation. Combination of two or more drugs in therapy may result in one of three outcomes: (1) additive, i.e., the effect of the combination is be equal to the sum of the effects of each drug when administered alone; (2) synergistic, i.e., the effect of the combination is greater than the sum of the effects of each drug when administered alone; or (3) antagonistic, i.e., the effect of the combination is less than the sum of the effects of each drug when administered alone. In one embodiment of the present invention, a subject is first treated with a substantially sub-toxic dose of a protein kinase inhibitor. After waiting for a period of time sufficient to allow development of a substantially efficacious response to the administration of the protein kinase inhibitor, a synergistic dose of a benzoquinone ansamycin is administered. Using this dosing schedule, a synergistic rather than additive effect of the two compounds is achieved. In one embodiment of the invention, the protein kinase inhibitor is a compound

listed in Table 2, and the benzoquinone ansamycin is a compound having formula (I), 17-AAG, or 17-DMAG. In another embodiment of the invention, the protein kinase inhibitor is a drug approved by the Federal Drug Administration as a stand-alone treatment for cancer, and the benzoquinone ansamycin is a compound having formula (I), 17-AAG, or 17-DMAG. In one preferred embodiment, the cytotoxic agent is Iressa and the benzoquinone ansamycin is 17-AAG or 17-DMAG.

[000100] In another embodiment of the invention, a subject is first treated with a first sub-toxic dose of a protein kinase inhibitor. After waiting for a period of time sufficient to allow development of a substantially efficacious response of the protein kinase inhibitor, a formulation comprising a synergistic dose of a benzoquinone ansamycin together with a second sub-toxic dose of the protein kinase inhibitor is administered. In general, the appropriate period of time sufficient to allow development of a substantially efficacious response to the protein kinase inhibitor will depend upon the pharmacokinetics of the protein kinase inhibitor, and will have been determined during clinical trials of therapy using the protein kinase inhibitor alone. In one embodiment of the invention, the period of time sufficient to allow development of a substantially efficacious response to the protein kinase inhibitor is between 1 hour and 96 hours. In another embodiment of the invention, the period of time sufficient to allow development of a substantially efficacious response to the protein kinase inhibitor is between 2 hours and 48 hours. In another embodiment of the invention, the period of time sufficient to allow development of a substantially efficacious response to the protein kinase inhibitor is between 4 hours and 24 hours.

[000101] The protein kinase inhibitors are selected from but are not limited to those listed in Table 2. As demonstrated below in Example 4 and in Figure 3A, pretreatment of cultured SKBr3 cells with the EGFR inhibitor Iressa followed by treatment with 17-AAG results in synergistic enhancement of the effects of Iressa. In contrast, the reverse order of administration or simultaneous administration results in an additive. Similar results were obtained with Iressa and 17-DMAG, as shown in Figure 3B. Thus to obtain the optimal synergistic effect, it is necessary to provide the protein kinase inhibitor first, wait a period of time sufficient to allow

development of a substantially efficacious response to the protein kinase inhibitor, and then provide the benzoquinone ansamycin.

[000102] In another embodiment, the subject is first treated with a sub-toxic dose of a benzoquinone ansamycin. After waiting for a period of time sufficient to allow development of a substantially efficacious response to the benzoquinone ansamycin, a synergistic dose of a microtubule stabilizing agent is administered. Using this dosing schedule, a synergistic rather than additive effect of the two compounds is achieved. Unexpectedly in light of Münster *et al.* "Modulation of Hsp90 function by ansamycins sensitizes breast cancer cells to chemotherapy-induced apoptosis in an RB- and schedule-dependent manner," *Clinical Cancer Research* (2001) 7: 2228-2236, pretreatment of cultured SKBr3 cells with the microtubule stabilizing agent paclitaxel followed by treatment with 17-AAG results in an additive effect, as demonstrated below in Example 4 and in Figure 5. In contrast, the reverse order of administration results in a synergistic effect.

Illustrative examples of microtubule stabilizing agents include but are not limited to paclitaxel, docetaxel, epothilone, discodermolide, and laulimalide. The benzoquinone ansamycin may be a compound having formula (I), 17-AAG, or 17-DMAG. In one embodiment of the invention, the microtubule stabilizing agent is paclitaxel an epothilone, discodermolide or an analog, or laulimalide or an analog. In a preferred embodiment of the invention, the microtubule stabilizing agent is epothilone D.

[000103] In another embodiment of the invention, the combination therapy the second agent is a drug approved by the Federal Drug Administration as a stand-alone treatment for cancer, and the benzoquinone ansamycin is a compound having formula (I) or is 17-AAG or 17-DMAG. Illustrative examples of suitable drugs include but are not limited to 5-fluorouracil, methotrexate, vinblastine, cyclophosphamide, mechlorethamine, chlorambucil, Melphalan, Ifosfamide, bleomycin, mitomycin and doxorubicin.

[000104] In another embodiment, the combination therapy may include an agent or procedure to mitigate potential side effects from the combination therapy agents. Diarrhea may be treated with antidiarrheal agents such as opioids (e.g. codeine, diphenoxylate, difenoxin, and loeramide), bismuth subsalicylate, and

octreotide. Nausea and vomiting may be treated with antiemetic agents such as dexamethasone, metoclopramide, diphenhydramine, lorazepam, ondansetron, prochlorperazine, thiethylperazine, and dronabinol. For those compositions that includes polyethoxylated castor oil such as Cremophor®, pretreatment with corticosteroids such as dexamethasone and methylprednisolone and/or H<sub>1</sub> antagonists such as diphenhydramine HCl and/or H<sub>2</sub> antagonists may be used to mitigate anaphylaxis.

[000105] The dose of the second agent when used in combination therapy with a benzoquinone ansamycin is determined based on the maximum tolerated dose observed when the second agent is used as the sole therapeutic agent (the “MTD”). In one embodiment of the invention, the dose of the second agent when used in combination therapy with a benzoquinone ansamycin is the MTD. In another embodiment of the invention, the dose of the second agent when used in combination therapy with a benzoquinone ansamycin is between 1% of the MTD and the MTD. In another embodiment of the invention, the dose of the second agent when used in combination therapy with a benzoquinone ansamycin is between 5% of the MTD and the MTD. In another embodiment of the invention, the dose of the second agent when used in combination therapy with a benzoquinone ansamycin is between 5% of the MTD and 75% of the MTD. In another embodiment of the invention, the dose of the second agent when used in combination therapy with a benzoquinone ansamycin is between 25% of the MTD and 75% of the MTD.

[000106] Use of the benzoquinone ansamycin allows for use of a lower therapeutic dose of the second agent, thus significantly widening the therapeutic window for treatment. In one embodiment, the therapeutic dose of the second agent is lowered by at least 10%. In another embodiment, the therapeutic dose of the second agent is lowered from 10 to 20%. In another embodiment, the therapeutic dose of the second agent is lowered from 20 to 50%. In another embodiment, the therapeutic dose of the second agent is lowered from 50 to 200%. In another embodiment, the therapeutic dose of the second agent is lowered from 100 to 1000%.

[000107] The MTD for a compound is determined using methods and materials known in the medical and pharmacological arts, for example through dose-

escalation experiments. One or more patients is first treated with a low dose of the compound, typically 10% of the dose anticipated to be therapeutic based on results of *in vitro* cell culture experiments. The patients are observed for a period of time to determine the occurrence of toxicity. Toxicity is typically evidenced as the observation of one or more of the following symptoms: vomiting, diarrhea, peripheral neuropathy, ataxia, neutropenia, or elevation of liver enzymes. If no toxicity is observed, the dose is increased 2-fold, and the patients are again observed for evidence of toxicity. This cycle is repeated until a dose producing evidence of toxicity is reached. The dose immediately preceding the onset of unacceptable toxicity is taken as the MTD.

[000108] The synergistic dose of the benzoquinone ansamycin used in combination therapy is determined based on the maximum tolerated dose observed when the benzoquinone ansamycin is used as the sole therapeutic agent. Clinical trials have determined an MTD for 17-AAG of 40 mg/m<sup>2</sup>. In one embodiment of the invention, the dose of the benzoquinone ansamycin when used in combination therapy is the MTD. In another embodiment of the invention, the dose of the benzoquinone ansamycin when used in combination therapy is between 1% of the MTD and the MTD. In another embodiment of the invention, the dose of the benzoquinone ansamycin when used in combination therapy is between 5% of the MTD and the MTD. In another embodiment of the invention, the dose of the benzoquinone ansamycin when used in combination therapy is between 5% of the MTD and 75% of the MTD. In another embodiment of the invention, the dose of the benzoquinone ansamycin when used in combination therapy is between 25% of the MTD and 75% of the MTD.

[000109] The dosages of the benzoquinone ansamycin and the Hsp90 client protein inhibitor when used in combination therapy may require further optimization depending upon the compounds being used, the disease or condition being treated, and the individual medical condition of the patient. Relevant factors include the activity of the specific compound employed; the age, body weight, general health, sex, and diet of the subject; the time and route of administration and the rate of excretion of the drug; and the severity of the condition being treated.

[000110] The present invention provides compositions of matter that are formulations of one or more active drugs and a pharmaceutically acceptable carrier. In one embodiment, the formulation comprises a novel benzoquinone ansamycin analog of the invention. In another embodiment, the formulation comprises a novel benzoquinone ansamycin analog of the invention as a mixture with an Hsp90 client protein inhibitor for use in combination therapy in accord with the methods of the present invention. In both of these embodiments, the active compounds may be in their free form or where appropriate as pharmaceutically acceptable derivatives such as prodrugs, esters, or salts.

[000111] The composition may be in any suitable form such as solid, semisolid, or liquid form. See Pharmaceutical Dosage Forms and Drug Delivery Systems, 5<sup>th</sup> edition, Lippicott Williams & Wilkins (1991), incorporated herein by reference. In general, the pharmaceutical preparation will contain one or more of the compounds of the invention as an active ingredient in admixture with an organic or inorganic carrier or excipient suitable for external, enteral, or parenteral application. The active ingredient may be compounded, for example, with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, suppositories, pessaries, solutions, emulsions, suspensions, and any other form suitable for use. The carriers that can be used include water, glucose, lactose, gum acacia, gelatin, mannitol, starch paste, magnesium trisilicate, talc, corn starch, keratin, colloidal silica, potato starch, urea, and other carriers suitable for use in manufacturing preparations, in solid, semi-solid, or liquified form. In addition, auxiliary stabilizing, thickening, and coloring agents and perfumes may be used.

[000112] In one embodiment, the compositions containing a compound useful in the methods of the invention are Cremophore®-free. Cremophore® (BASF Aktiengesellschaft) is a polyethoxylated castor oil which is typically used as a surfactant in formulating low soluble drugs. However, because Cremophore® can cause allergic reactions in a subject, compositions that minimize or eliminate Cremophore® are preferred.

[000113] Where applicable, the compounds useful in the methods of the invention may be formulated as microcapsules and nanoparticles. General protocols are described for example, by Microcapsules and Nanoparticles in Medicine and

Pharmacy by Max Donbrow, ed., CRC Press (1992) and by U.S. Patent Nos. 5,510,118; 5,534,270; and 5,662,883 which are all incorporated herein by reference. By increasing the ratio of surface area to volume, these formulations allow for the oral delivery of compounds that would not otherwise be amenable to oral delivery.

5           **[000114]** The compounds useful in the methods of the invention may also be formulated using other methods that have been previously used for low solubility drugs. For example, the compounds may form emulsions with vitamin E or a PEGylated derivative thereof as described by PCT publications WO 98/30205 and WO 00/71163, each of which is incorporated herein by reference. Typically, the  
10       compound useful in the methods of the invention is dissolved in an aqueous solution containing ethanol (preferably less than 1% w/v). Vitamin E or a PEGylated-vitamin E is added. The ethanol is then removed to form a pre-emulsion that can be formulated for intravenous or oral routes of administration. Another method involves encapsulating the compounds useful in the methods of the invention in  
15       liposomes. Methods for forming liposomes as drug delivery vehicles are well known in the art. Suitable protocols include those described by U.S. Patent Nos. 5,683,715; 5,415,869, and 5,424,073 which are incorporated herein by reference relating to another relatively low solubility cancer drug paclitaxel and by PCT  
20       Publication WO 01/10412 which is incorporated herein by reference relating to epothilone B. Of the various lipids that may be used, particularly preferred lipids for making encapsulated liposomes include phosphatidylcholine and polyethyleneglycol-derivitized distearyl phosphatidyl-ethanolamine.

**[000115]** Yet another method involves formulating the compounds useful in the methods of the invention using polymers such as polymers such as biopolymers  
25       or biocompatible (synthetic or naturally occurring) polymers. Biocompatible polymers can be categorized as biodegradable and non-biodegradable. Biodegradable polymers degrade *in vivo* as a function of chemical composition, method of manufacture, and implant structure. Illustrative examples of synthetic polymers include polyanhydrides, polyhydroxyacids such as polylactic acid,  
30       polyglycolic acids and copolymers thereof, polyesters polyamides polyorthoesters and some polyphosphazenes. Illustrative examples of naturally occurring polymers

include proteins and polysaccharides such as collagen, hyaluronic acid, albumin, and gelatin.

5 [000116] Another method involves conjugating the compounds useful in the methods of the invention to a polymer that enhances aqueous solubility. Examples of suitable polymers include polyethylene glycol, poly-(d-glutamic acid), poly-(l-glutamic acid), poly-(l-glutamic acid), poly-(d-aspartic acid), poly-(l-aspartic acid), poly-(l-aspartic acid) and copolymers thereof. Polyglutamic acids having molecular weights between about 5,000 to about 100,000 are preferred, with molecular weights between about 20,000 and 80,000 being more preferred and with molecular weights between about 30,000 and 60,000 being most preferred. The polymer is conjugated via an ester linkage to one or more hydroxyls of an inventive geldanamycin using a protocol as essentially described by U.S. Patent No. 5,977,163 which is incorporated herein by reference.

15 [000117] In another method, the compounds useful in the methods of the invention are conjugated to a monoclonal antibody. This method allows the targeting of the inventive compounds to specific targets. General protocols for the design and use of conjugated antibodies are described in Monoclonal Antibody-Based Therapy of Cancer by Michael L. Grossbard, ed. (1998), which is incorporated herein by reference.

20 [000118] The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the subject treated and the particular mode of administration. For example, a formulation for intravenous use comprises an amount of the inventive compound ranging from about 1 mg/mL to about 25 mg/mL, preferably from about 5 mg/mL to 15 mg/mL, and more preferably about 10 mg/mL. Intravenous formulations are typically diluted between about 2 fold and about 30 fold with normal saline or 5% dextrose solution prior to use.

30 [000119] In one aspect of the present invention, the compounds useful in the methods of the invention are used to treat cancer. In one embodiment, the compounds of the present invention are used to treat cancers of the head and neck which include but are not limited to tumors of the nasal cavity, paranasal sinuses, nasopharynx, oral cavity, oropharynx, larynx, hypopharynx, salivary glands, and



paragangliomas. In another embodiment, the compounds of the present invention are used to treat cancers of the liver and biliary tree, particularly hepatocellular carcinoma. In another embodiment, the compounds of the present invention are used to treat intestinal cancers, particularly colorectal cancer. In another  
5 embodiment, the compounds of the present invention are used to treat ovarian cancer. In another embodiment, the compounds of the present invention are used to treat small cell and non-small cell lung cancer. In another embodiment, the compounds of the present invention are used to treat breast cancer. In another embodiment, the compounds of the present invention are used to treat sarcomas,  
10 including fibrosarcoma, malignant fibrous histiocytoma, embryonal rhabdomyosarcoma, leiomyosarcoma, neuro-fibrosarcoma, osteosarcoma, synovial sarcoma, liposarcoma, and alveolar soft part sarcoma. In another embodiment, the compounds of the present invention are used to treat neoplasms of the central nervous systems, particularly brain cancer. In another embodiment, the  
15 compounds of the present invention are used to treat lymphomas which include Hodgkin's lymphoma, lymphoplasmacytoid lymphoma, follicular lymphoma, mucosa-associated lymphoid tissue lymphoma, mantle cell lymphoma, B-lineage large cell lymphoma, Burkitt's lymphoma, and T-cell anaplastic large cell lymphoma.

20 [000120] In another embodiment, the compounds and compositions useful in the methods of the invention are used at sub-cytotoxic levels in combination with other agents in order to achieve highly selective activity in the treatment of non-cancerous diseases. In this embodiment, the compounds useful in the methods of the invention are used to reduce the cellular levels of Hsp90 client proteins, which are  
25 then effectively inhibited by the second agent. Binding of the client proteins to Hsp90 stabilizes the client proteins and maintains them in a soluble, inactive form ready to respond to activating stimuli. Binding of a benzoquinone ansamycin analog to Hsp90 results in targeting of the client protein to the proteasome, and subsequent degradation. For systems such as the steroid receptor, however, Hsp90 forms an  
30 integral part of the functional receptor complex along with several other proteins such as Hsp70, Hsp40, p23, hip, Hsp56, and immunophilins. Hsp90 appears to regulate the activity of the steroid receptor by maintaining the receptor in a high-

affinity hormone-binding conformation. Binding of geldanamycin to Hsp90 appears to result in dissociation of p23 from the complex and reduce the level of hormone binding to the receptor (Fliss *et al.* (2000) "Control of estrogen receptor ligand binding by Hsp90," *J. Steroid Biochem. Mol. Biol.* 75: 223-30; Kimmins and MacRae (2000), "Maturation of steroid receptors: an example of functional cooperation among molecular chaperones and their associated proteins," *Cell Stress Chaperones* 5: 76-86.

[000121] Thus, Hsp90 inhibitors such as geldanamycin, geldanamycin analogs, radicicol, and the like can be used in accord with the methods of the present invention to alter the function of hormone receptors, making it easier to inhibit the associated signal pathways using low levels of a second drug which targets the proteins involved in those signaling pathways. Such a combination therapy can be useful to reduce non-specific toxicity associated with therapy by reducing the levels of the drugs required.

[000122] In another embodiment, the compounds useful in the methods of the invention are used to treat non-cancerous diseases or conditions characterized by undesired cellular hyperproliferation, including neurodegenerative diseases, psoriasis, stenosis, and restenosis.

[000123] In another embodiment, the compounds useful in the methods of the invention are used in combination with other agents as described above to treat non-cancerous diseases or conditions characterized by undesired cellular hyperproliferation, including neuro-degenerative diseases, psoriasis, stenosis, and restenosis. Specific examples of non-cancerous diseases treatable by this combination therapy include neurodegenerative diseases, such as Alzheimer's, Parkinson's, Huntington's and the like. There is evidence that the immunophilin FKBP-52 is involved as a client protein for Hsp90 in the formation of various steroid receptor complexes and plays a role in the regeneration of damaged neurons (Gold *et al.*, "Immunophilin FK506-Binding Protein 52 (Not FK506-Binding Protein 12) Mediates the Neurotrophic Action of FK506," 1999, *J. Pharmacology & Exp. Ther.* 289: 1202-1210). The combination of geldanamycin, a geldanamycin analog, radicicol, or the like with a second agent which binds to and/or inhibits

FKBP-52 in accord with the methods of the present invention can thus be used to treat neurodegenerative diseases.

[000124] Further examples of non-cancerous diseases treatable by the combination therapy of the present invention include non-cancerous diseases characterized by cellular hyperproliferation, such as psoriasis, stenosis, and restenosis. Cell proliferation is regulated by protein tyrosine kinases, many of which are known to be client proteins for Hsp90. Psoriasis is thought to involve the epidermal growth factor receptor (EGFR), a protein tyrosine kinase, and inhibitors of EGFR have been proposed as treatments for psoriasis (Ben-Bassat & Klein, "Inhibitors of Tyrosine Kinases in the Treatment of Psoriasis," (2000), *Curr. Pharm. Des.* 6: 933-942). Geldanamycin and herbimycin have been shown to block maturation of EGFR (Sakagami *et al.*, "Benzoquinoid ansamycins (herbimycin A and geldanamycin) interfere with the maturation of growth factor receptor tyrosine kinases," (1999) *Cell Stress Chaperones* 4: 19-28). The combination of geldanamycin or a geldanamycin analog with an inhibitor of EGFR in accord with the methods of the present invention can thus be used to treat psoriasis.

[000125] The compounds useful in the methods of the invention may also be used to treat stenosis and restenosis, particularly associated with *in vivo* devices such as stents. In one embodiment, the compounds useful in the methods of the invention are used to coat stents and other surgically-implantable devices. In another embodiment, the compounds useful in the methods of the invention are used in combination with other agents as described above to coat stents, catheters, prostheses, and other *in vivo* devices.

[000126] A detailed description of the invention having been provided above, the following examples are given for the purpose of illustrating the present invention and shall not be construed as being a limitation on the scope of the invention or claims.

### EXAMPLE 1

#### Preparation of 17-alkylamino-17-desmethoxygeldanamycins

[000127] The general procedure for preparing 17-alkylamino-17-desmethoxygeldanamycins is as follows. The geldanamycin analog (25 mmol) is suspended in 350 mL of anhydrous  $\text{CH}_2\text{Cl}_2$  under inert atmosphere. A solution of the alkylamine (50 mmol) in 10 mL of  $\text{CH}_2\text{Cl}_2$  is added dropwise. After 1 hour, the mixture is evaporated to dryness and the residue is dissolved in 50 mL of chloroform and precipitated by addition of 600 mL of hexane. Filtration and vacuum drying yields the product. The products are characterized by NMR and LC/MS.

Compounds prepared according to this method include:

- 17-allylamino-17-desmethoxygeldanamycin;
- 17-(2-(dimethylamino)ethyl)amino-17-desmethoxygeldanamycin;
- 17-ethylamino-17-desmethoxygeldanamycin;
- 17-propylamino-17-desmethoxygeldanamycin;
- 17-butylamino-17-desmethoxygeldanamycin;
- 17-(cyclopropyl)methylamino-17-desmethoxygeldanamycin;
- 17-cyclobutylamino-17-desmethoxygeldanamycin;
- 17-(2-phenylcyclopropyl)amino-17-desmethoxygeldanamycin;
- 17-(2-fluoroethyl)amino-17-desmethoxygeldanamycin;
- 17-(2,2-difluoroethyl)amino-17-desmethoxygeldanamycin;
- 17-azetidiny-17-desmethoxygeldanamycin;
- 17-amino-17-desmethoxygeldanamycin;
- 17-(3-(dimethylamino)propyl)amino-17-desmethoxygeldanamycin;
- 17-(4-(dimethylamino)butyl)amino-17-desmethoxygeldanamycin;
- 17-(3-dimethylamino)-2-propylamino-17-desmethoxygeldanamycin;
- 17-(2-dimethylamino)-1-propylamino-17-desmethoxygeldanamycin;
- 17-(N-ethylpyrrolidin-2-yl)methylamino-17-desmethoxygeldanamycin;
- 17-(1-pyrazinyl)ethylamino-17-desmethoxygeldanamycin;
- 17-(2-bis(2-hydroxyethyl)amino)ethylamino-17-desmethoxygeldanamycin;
- 17-(N-methylpyrrolidin-2-yl)ethylamino-17-desmethoxygeldanamycin;
- 17-(3-(diethylamino)propyl)amino-17-desmethoxygeldanamycin;
- 17-(3-(4-morpholino)propyl)amino-17-desmethoxygeldanamycin;
- 17-(4-imidazolyl)ethylamino-17-desmethoxygeldanamycin;
- 17-(1-methyl-4-imidazolyl)ethylamino-17-desmethoxygeldanamycin;

17-(1-methyl-5-imidazolyl)ethylamino-17-desmethoxygeldanamycin;  
17-(4-pyridyl)ethylamino-17-desmethoxygeldanamycin;  
17-(2-hydroxyethyl)amino-17-desmethoxygeldanamycin;  
17-(1-hydroxymethyl)propylamino-17-desmethoxygeldanamycin;  
5 17-(2-(hydroxymethyl)cyclohexyl)amino-17-desmethoxygeldanamycin;  
17-(2-dioxolan-1-yl)ethylamino-17-desmethoxygeldanamycin;  
17-(3,3-dimethoxypropyl)amino-17-desmethoxygeldanamycin;  
17-(2-*tert*-butoxycarbonylamino)ethylamino-17-desmethoxygeldanamycin; and  
17-(*N*-methyl 2-*tert*-butoxycarbonylamino)ethylamino-17-desmethoxygeldanamycin.

10

### EXAMPLE 2

#### Preparation of 11-O-allyl-17-alkylamino-17-desmethoxygeldanamycins

[000128] The general procedure for preparing 11-O-allyl-17-alkylamino-17-  
15 desmethoxy-geldanamycins is as follows. The 17-alkylamino-17-desmethoxy-  
geldanamycin analog (10 mmol), allyl *tert*-butyl carbonate (14 mmol), palladium  
acetate (0.1 mmol), and triphenylphosphine (0.67 mmol) are dissolved in 60 mL of  
freshly distilled, air-free tetrahydrofuran under inert atmosphere. The mixture is  
heated at reflux for 16 hours, and is monitored by thin-layer chromatography. Upon  
20 completion, the solvents are removed in vacuo and the product isolated by silica gel  
chromatography.

### EXAMPLE 3

#### Preparation of 11,17-linked-17-amino-17-desmethoxygeldanamycins

25

[000129] The general procedure for preparing 11,17-linked-17-amino-17-  
desmethoxy-geldanamycins is as follows. The 11-O-allyl-17-  
alkylaminogeldanamycin analog (10 mmol), and benzylidene-  
bis(tricyclohexylphosphine)ruthenium dichloride (5 mmol) are dissolved in 250 mL  
30 of anhydrous CH<sub>2</sub>Cl<sub>2</sub> under inert atmosphere, and the reaction is monitored by thin-  
layer chromatography. After 24 hours, the reaction is concentrated and the residue  
is purified by silica gel chromatography.

**EXAMPLE 4****Interaction of Geldanamycin Analogs and Cytotoxic Agents in SKBr3 Cells**

5           [000130] SKBr3 and H358 cell lines were obtained from American Type Culture Collection (Manassas, VA). Cells were maintained in McCoy's 5A medium with 10% fetal bovine serum. 17-AAG and 17-DMAG were dissolved in DMSO at 10 mM and stored at -20°C.

10           [000131] Cells were seeded in duplicate in opaque-walled 96-well microtiter plates at 4000 cells per well and allowed to attach overnight. Serial dilutions of each drug were added, and the cells were incubated for 72 hours. The IC<sub>50</sub> was determined using the CellTiter-Glo Luminescent Cell Viability Assay (Promega, Madison, WI), which correlates with the number of live cells.

15           [000132] For the drug combination assay, cells were seeded in duplicate in 96-well plates (4000 cells/well). After an overnight incubation, cells were treated with drug alone or in combination. Based on the IC<sub>50</sub> values of each individual drug, combined drug treatment was designed at constant ratios of two drugs, i.e., equivalent to the ratio of their IC<sub>50</sub>. Three different treatment schedules were used. The first treatment schedule used simultaneous exposure to both 17-AAG and the  
20           second cytotoxic agent for 72 hours. In the second schedule, the cells were exposed to 24 hours of 17-AAG or 17-DMAG. The second cytotoxic agent was then added to the cells and incubated for 48 hours. In the third treatment schedule, cells were exposed to the second cytotoxic agent alone for 24 hours followed by addition of 17-AAG or 17-DMAG for 48 hours. Cell viability was determined by luminescent  
25           assay (Promega). Combination analysis was performed using CalcuSyn software (Biosoft, Cambridge, UK). The "combination index" (CI) refers to a measure of the additivity of the effects of two drugs administered in combination, and was calculated according to the formula:

$$CI = [D]_1/[D_x]_1 + [D]_2/[D_x]_2$$

30           wherein [D]<sub>1</sub> and [D]<sub>2</sub> represent the concentrations of the first and second drug, respectively, that in combination provide a response of x% in the assay, and [D<sub>x</sub>]<sub>1</sub> and [D<sub>x</sub>]<sub>2</sub> represent the concentrations of the first and second drug, respectively, that when

used alone provide a response of x% in the assay. According to this formula, a CI = 1 indicates a simple additive effect of the two drugs in combination, whereas a CI < 1 indicates synergism and a CI > 1 indicates antagonism between the two drugs.

5 [000133] Addition of 17-AAG or 17-DMAG to cells after exposure to Iressa synergistically increased the cytotoxicity of Iressa (Figure 3). An additive effect was detected when cells were exposed to 17-AAG or 17-DMAG before Iressa. Exposure of cells to 17-AAG or 17-DMAG and Iressa simultaneously also produced an additive effect. Thus the optimal synergistic effect is only observed when there is prior exposure of the cells to the protein kinase inhibitor, followed by exposure to  
10 the benzoquinone ansamycin after waiting for a period of time to allow development of efficacy of the protein kinase inhibitor.

[000134] In contrast, addition of 17-AAG or 17-DMAG to cells after exposure to paclitaxel additively increased the cytotoxicity of paclitaxel (Figure 5). A synergistic effect was observed when cells were exposed to paclitaxel first, and  
15 were later treated with 17-AAG or 17-DMAG. In this case, optimal synergism is only observed when the cells are first exposed to the benzoquinone ansamycin, followed by exposure to the microtubule stabilizing agent after waiting for a period of time to allow development of efficacy of the benzoquinone ansamycin.

20

### EXAMPLE 5

#### Production of 28-desmethylgeldanamycin by AT domain substitution

[000135] Substitution of the acyltransferase domain in module 1 of the  
25 geldanamycin PKS gene with an AT domain specific for malonyl-CoA instead of 2-methylmalonyl-CoA results in formation of 28-desmethyl-geldanamycin. The domain substitution is created by introducing a malonyl-CoA specific acyltransferase domain from a heterologous PKS gene, for example from the rapamycin, tylosin, or FK520 PKS genes or the like, into the geldanamycin PKS  
30 locus by homologous recombination into a geldanamycin producing strain, aided by a selectable antibiotic resistance gene, then isolating the recombinants resulting from double crossover events in which the wild-type acyltransferase domain is replaced

with one specific for malonyl-CoA. The AT domain of module 1 is encoded by nucleotides 1626 through 2670, approximately, of SEQ ID NO:1. This sequence information together with the methods described in U.S. patents 6,399,789; 6,403,775; and 5,962,290 allows one skilled in the art to construct recombination  
5 vectors that result in replacement of the native AT domain of module 1 with an AT domain having a specificity for malonyl-CoA. Suitable examples of AT domains with specificity for malonyl-CoA may be found in the rapamycin PKS genes (modules 2, 5, 8, 9, 11, 12, and 14), as described in U.S. patent 6,399,789, as well as the tylosin PKS genes (modules 3 and 7) as described in U.S. patent 5,876,991; the spiramycin genes (modules 1-3 and 7), as described in U.S. Patent 5,945,320; the  
10 FK520 genes (modules 3 and 10), as described in WO 00/20601; the pikromycin genes (module 2) as described in WO 99/61599; the narbomycin genes (module 2), as described in U.S. patent 6,303,767; the avermectin genes (module 2), and others. Each of the above documents is incorporated herein by reference. Fermentation of a  
15 host cell comprising the resulting hybrid PKS together with the remaining geldanamycin biosynthetic genes under conditions wherein the native strain produces geldanamycin, followed by extraction of the broth and purification provides 28-desmethyl-geldanamycin.

#### EXAMPLE 6

##### Production of 28-desmethylgeldanamycin by site-directed mutagenesis

c14-me  
20  
[000136] The acyltransferase domain in module 1 of the geldanamycin PKS gene is mutagenized according to the methods described in Reeves *et al.*, "Alteration  
25 of the substrate specificity of a modular polyketide synthase acyltransferase domain through site-directed mutagenesis," *Biochemistry* 2001, 40: 15464-15470, and in U.S. patent application serial no. 60/310,730, entitled "Alteration of the substrate specificity of a modular PKS AT domain," which is incorporated herein by reference. The amino acid sequence Tyr-Ala-Ser-His (SEQ ID NO: 3), encoded by  
30 nucleotide sequence TAC-GCC-TCC-CAC (SEQ ID NO: 4) at positions 2194 to 2205 in SEQ ID NO:1, is mutagenized using methods known to one skilled in the art to generate the mutant amino acid sequence His-Ala-Phe-His (SEQ ID NO: 5), for



example by mutagenesis of the nucleotide sequence to CAC-GCC-TTC-CAC (SEQ ID NO: 6) as described in the Reeves *et al.* reference cited above. Fermentation of a host cell comprising the resulting mutagenized PKS together with the remaining geldanamycin biosynthetic genes under conditions wherein the native strain produces geldanamycin, followed by extraction of the broth and purification provides 28-desmethyl-geldanamycin.

#### EXAMPLE 7

##### Production of 4,5-dihydro-5-hydroxygeldanamycin by reductive domain swap

[000137] The coding sequence for the reduction cassette of module 6, which has both DH and KR domains, is replaced with a coding sequence for a reduction cassette that has only a KR domain. The reduction cassette is contained in the sequence between the end of the AT domain, at approximately nucleotide position 2805 of SEQ ID NO:2, and the beginning of the ACP domain, at approximately nucleotide position 6028 of SEQ ID NO:2. This sequence information together with the methods described in U.S. patents 6,399,789; 6,403,775; and 5,962,290 allows one skilled in the art to construct recombination vectors that result in replacement of the native reduction cassette of module 6 with a cassette encoding only a KR domain. Suitable examples of cassettes encoding only a KR domain may be found in the erythromycin and rapamycin PKS genes, as described in U.S. patent 6,399,789. Fermentation of a host cell comprising the resulting hybrid PKS together with the remaining geldanamycin biosynthetic genes under conditions wherein the native strain produces geldanamycin, followed by extraction of the broth and purification provides 4,5-dihydro-5-hydroxy-geldanamycin.

#### EXAMPLE 8

##### Production of 4,5-dihydro-5-hydroxygeldanamycin by site-directed mutagenesis

[000138] Inactivation of the dehydratase domain in module 6 of the geldanamycin PKS gene by site-specific mutation of the wild-type domain results in

production of 4,5-dihydro-5-hydroxygeldanamycin. The DH domain of module 6 is encoded by nucleotides 2805 to 4276, approximately, of SEQ ID NO:2. Two particular sequences may be targeted for mutational inactivation of the DH domain. In one embodiment, the DNA sequence encoding the DH peptide motif His-Val-Ile-Ser-Gly-Ala-Val-Leu-Val-Pro (SEQ ID NO: 7), nucleotides 2956 through 2985 of SEQ ID NO:2, is mutated so as to produce a peptide having an amino acid other than histidine at the first position. The CAC codon encoding histidine is mutated, for example to CAA or CAG to encode a glutamine, as illustrated in SEQ ID NO: 8. Fermentation of a host cell comprising the resulting mutagenized PKS together with the remaining geldanamycin biosynthetic genes under conditions wherein the native strain produces geldanamycin, followed by extraction of the broth and purification provides 4,5-dihydro-5-hydroxy-geldanamycin.

#### EXAMPLE 9

##### Production of 4,5-dihydro-5-hydroxygeldanamycin by deletion

[000139] A portion of the nucleotide sequence in module 6 between the end of the AT domain (approximately nucleotide 2805 of SEQ ID NO:2) and the start of the KR domain (approximately nucleotide 5212 of SEQ ID NO:2) is deleted to provide a modified PKS for production of 4,5-dihydro-5-hydroxy-geldanamycin. The nucleotide sequence between 2805 and 3270, approximately, of SEQ ID NO:2 is deleted so as to leave a linker region between the AT and KR domains of approximately 465 amino acids. Fermentation of a host cell comprising the resulting modified PKS together with the remaining geldanamycin biosynthetic genes under conditions wherein the native strain produces geldanamycin, followed by extraction of the broth and purification provides 4,5-dihydro-5-hydroxy-geldanamycin.

#### EXAMPLE 10

##### Production of 15-hydroxygeldanamycin by reductive domain swap

[000140] The reduction cassette in module 1 is encoded by the sequence between the end of the AT domain, at approximately nucleotide position 2670 of

SEQ ID NO:1, and the beginning of the ACP domain, at approximately nucleotide position 5895 of SEQ ID NO:1. This sequence information together with the methods described in U.S. patents 6,399,789; 6,403,775; and 5,962,290 allows one skilled in the art to construct recombination vectors that result in replacement of the native reduction cassette of module 1 with a cassette encoding only a KR domain. Suitable examples of cassettes encoding only a KR domain may be found in the erythromycin and rapamycin PKS genes, as described in U.S. patent 6,399,789. Fermentation of a host cell comprising the resulting hybrid PKS together with the remaining geldanamycin biosynthetic genes under conditions wherein the native strain produces geldanamycin, followed by extraction of the broth and purification provides 15-hydroxy-geldanamycin.

#### EXAMPLE 11

##### Production of 15-hydroxygeldanamycin by site-directed mutagenesis

[000141] Inactivation of the dehydratase domain in module 1 of the geldanamycin PKS gene by site-specific mutation of the wild-type domain results in production of 15-hydroxygeldanamycin. The DH domain of module 1 is encoded by nucleotides 2670 to 4140, approximately, of SEQ ID NO:1. Two particular sequences may be targeted for mutational inactivation of the DH domain. In one embodiment, the DNA sequence encoding the DH peptide motif His-Ala-Val-Ser-Gly-Thr-Val-Leu-Leu-Pro (SEQ ID NO: 9), nucleotides 2821 through 2850 of SEQ ID NO:1, is mutated so as to produce a peptide having an amino acid other than histidine at the first position. The CAC codon encoding histidine is mutated, for example to CAA or CAG to encode a glutamine as illustrated in SEQ ID NO: 10. Fermentation of a host cell comprising the resulting mutagenized PKS together with the remaining geldanamycin biosynthetic genes under conditions wherein the native strain produces geldanamycin, followed by extraction of the broth and purification provides 15-hydroxy-geldanamycin.

#### EXAMPLE 12

##### 8,9-epoxy-8,9-dihydrogeldanamycin

[000142] To a solution of geldanamycin (120 mg, 0.21 mmol) in dichloromethane (5 mL) was added 85% 3-chloroperoxybenzoic acid (93 mg, 0.42 mmol). After the mixture was stirred at room temperature overnight, TLC showed the reaction to be complete. The reaction mixture was diluted with ethyl acetate and washed subsequently with aqueous sodium sulfite, aqueous bicarbonate, and brine. The organic solution was dried over anhydrous sodium sulfate, filtered, and evaporated to dryness. The crude product was purified by flash chromatography on silica gel, giving 30 mg of 8,9-epoxy-8,9-dihydrogeldanamycin as a yellow solid. The compound was characterized by NMR and MS spectrometry.

### EXAMPLE 13

#### 8,9-epoxy -17-[2-(dimethylamino)ethylamino]-8,9-dihydro-17-demethoxygeldanamycin

[000143] To a solution of 8,9-epoxy-8,9-dihydrogeldanamycin (20 mg, 0.035 mmol) in 1,2-dichloroethane (2 mL) was added *N,N*-dimethylethylenediamine (10  $\mu$ L, 0.09 mmol). The mixture was stirred at room temperature overnight. The reaction was worked up by diluting with ethyl acetate and washing sequentially with aqueous bicarbonate and brine. The organic solution was dried over anhydrous sodium sulfate, filtered, and evaporated to dryness. The crude product was purified by HPLC on a C-18 column, giving 10 mg of 8,9-epoxy -17-[2-(dimethylamino)ethylamino]-8,9-dihydro-17-demethoxygeldanamycin as a purple solid. The compound was characterized by NMR and MS spectrometry.

### EXAMPLE 14

#### 17-[2-(dimethylamino)ethylamino]-17-demethoxygeldanamycin

[000144] To a solution of geldanamycin (60 mg, 0.1 mmol) in 1,2-dichloroethane (4 mL) was added *N,N*-dimethylethylenediamine (22  $\mu$ L, 0.2 mmol). The mixture was stirred at room temperature overnight. The reaction was diluted with ethyl acetate and washed sequentially with aqueous bicarbonate and brine. The organic solution was dried over anhydrous sodium sulfate, filtered, and evaporated

to dryness. The crude product was purified by flash chromatography on silica gel, giving 59 mg of 17-[2-(dimethylamino)-ethylamino]-17-demethoxygeldanamycin as a purple solid. The compound was characterized by NMR and MS spectrometry.

5

**EXAMPLE 15****17-[2-(dimethylamino)ethylamino]-11-oxo-17-demethoxygeldanamycin**

[000145] To a solution of 17-[2-(dimethylamino)ethylamino]-17-demethoxygeldanamycin (50 mg, 0.08 mmol) in chloroform (5 ml) was added the Dess-Martin periodinane (208 mg, 0.49 mmol). The mixture was heated to reflux for one hour. The reaction mixture was diluted in ethyl acetate and washed subsequently with aqueous sodium thiosulfate, aqueous bicarbonate, and brine. The organic solution was dried over anhydrous sodium sulfate, filtered, and evaporated to dryness. The crude product was purified by HPLC on a C-18 column, giving 26 mg of 17-[2-(dimethylamino)ethylamino]-11-oxo-17-demethoxygeldanamycin as a purple solid. The compound was characterized by NMR and MS spectrometry.

15

**EXAMPLE 16****17-[2-(dimethylamino)ethylamino]-11-oxo-17-demethoxygeldanamycin-11-oxime**

[000146] To a solution of 17-[2-(dimethylamino)ethylamino]-11-oxo-17-demethoxy-geldanamycin (30 mg, 0.049 mmol) in ethanol (4 mL) was added triethylamine (60  $\mu$ L, 0.43 mmol) and hydroxylamine hydrochloride (30 mg, 0.4 mmol). After the mixture was stirred at room temperature for 3 h, TLC showed reaction complete. After ethanol was evaporated, the residue was dissolved in chloroform and water. The organic layer was dried over anhydrous sodium sulfate, filtered, and evaporated to dryness. The crude product was purified by flash chromatography on silica gel, giving 10 mg of 17-[2-(dimethylamino)ethylamino]-11-oxo-17-demethoxygeldanamycin-11-oxime as a purple solid. The compound was characterized by NMR and MS spectrometry.


25

30

## CLAIMS

What is claimed is:

1. A compound or pharmaceutically acceptable salt or prodrug thereof having the formula:



- wherein  $R^1$  is MeO,  $(CH_2)_3N$  or  $R^9-NH$ , wherein  $R^9$  is selected from the group consisting of H, substituted or unsubstituted  $C_1-C_6$  alkyl, substituted or unsubstituted  $C_1-C_6$  alkenyl, substituted or unsubstituted  $C_1-C_6$  alkynyl, substituted or unsubstituted  $C_3-C_6$  cycloalkyl, piperidinyl, N-alkylpiperidinyl, hexahydropyranyl, furfuryl, tetrahydrofurfuryl, pyrrolidinyl, N-alkylpyrrolidinyl, piperazinylamino, N-alkylpiperazinyl, morpholinyl, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-methyl-pyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl, or  $R^6$  is H and  $R^1$  and  $R^5$  taken together form a group of the formula  $NH-Z-O$ , wherein Z is a linker comprised of from 1 to 6 carbon atoms and 0 to 2 nitrogen atoms and wherein the O is attached at the position of  $R^5$ ;
- $R^2$  is selected from the group consisting of H, halogen,  $OR^{10}$ ,  $NHR^{10}$ ,  $SR^{10}$ , aryl, and heteroaryl, wherein  $R^{10}$  is selected from the group consisting of substituted or unsubstituted  $C_1-C_6$  alkyl, substituted or unsubstituted  $C_1-C_6$  alkenyl, substituted or unsubstituted  $C_1-C_6$  alkynyl, and substituted or unsubstituted  $C_3-C_6$  cycloalkyl;
- $R^3$  is H, OH, or OMe;

R<sup>4</sup> is H or Me;

R<sup>5</sup> is OH or O-C(=O)-CH<sub>2</sub>NH<sub>2</sub> and R<sup>6</sup> is H, or R<sup>5</sup> and R<sup>6</sup> taken together form =O or =N-OR<sup>11</sup>, wherein R<sup>11</sup> is selected from the group consisting of H, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkenyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkynyl, substituted or unsubstituted C<sub>3</sub>-C<sub>6</sub> cycloalkyl, aryl, or heteroaryl;

R<sup>7</sup> is H and R<sup>8</sup> is H or OH, or R<sup>7</sup> and R<sup>8</sup> taken together form a bond; and X is O or a bond, with the provisos that when R<sup>3</sup> is H, R<sup>4</sup> is Me, and R<sup>7</sup> is H and R<sup>8</sup> is H or R<sup>7</sup> and R<sup>8</sup> taken together form a bond that either R<sup>6</sup> is H and R<sup>1</sup> and R<sup>5</sup> taken together form a group of the formula NH-Z-O, wherein Z is a linker comprised of from 1 to 6 carbon atoms and 0 to 2 nitrogen atoms and wherein the O is attached at the position of R<sup>5</sup>, or that R<sup>1</sup> is (CH<sub>2</sub>)<sub>3</sub>N or R<sup>9</sup>-NH, wherein R<sup>9</sup> is selected from the group consisting of piperidiny, N-alkylpiperidiny, hexahydropyrany, furfuryl, tetrahydrofurfuryl, pyrrolidiny, N-alkylpyrrolidiny, piperazinylamino, N-alkylpiperazinyl, morpholiny, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-methyl-pyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl; and that when R<sup>3</sup> is H and R<sup>4</sup> is Me that R<sup>7</sup> is H and R<sup>8</sup> is OH.

2. A compound of Claim 1 wherein R<sup>1</sup> is (CH<sub>2</sub>)<sub>3</sub>N or R<sup>9</sup>-NH, wherein R<sup>9</sup> is selected from the group consisting of piperidiny, N-alkylpiperidiny, hexahydropyrany, furfuryl, tetrahydrofurfuryl, pyrrolidiny, N-alkylpyrrolidiny, piperazinylamino, N-alkylpiperazinyl, morpholiny, N-alkylaziridinylmethyl, (1-azabicyclo[1.3.0]hex-1-yl)ethyl, 2-(N-methyl-pyrrolidin-2-yl)ethyl, 2-(4-imidazolyl)ethyl, 2-(1-methyl-4-imidazolyl)ethyl, 2-(1-methyl-5-imidazolyl)ethyl, 2-(4-pyridyl)ethyl, and 3-(4-morpholino)-1-propyl;

R<sup>2</sup> is selected from the group consisting of H, halogen, OR<sup>10</sup>, NHR<sup>10</sup>, SR<sup>10</sup>, aryl, and heteroaryl, wherein R<sup>10</sup> is selected from the group consisting of substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkenyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkynyl, and substituted or unsubstituted C<sub>3</sub>-C<sub>6</sub> cycloalkyl;

R<sup>3</sup> is H, OH, or OMe;

R<sup>4</sup> is H or Me;

R<sup>5</sup> is OH or O-C(=O)-CH<sub>2</sub>NH<sub>2</sub> and R<sup>6</sup> is H, or R<sup>5</sup> and R<sup>6</sup> taken together form =O or =N-OR<sup>11</sup>, wherein R<sup>11</sup> is selected from the group consisting of H, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkenyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkynyl, substituted or unsubstituted C<sub>3</sub>-C<sub>6</sub> cycloalkyl, aryl, or heteroaryl;

R<sup>7</sup> is H and R<sup>8</sup> is H or OH, or R<sup>7</sup> and R<sup>8</sup> taken together form a bond; and  
X is O or a bond.

3. A compound of Claim 1 wherein R<sup>3</sup> is OH or OMe.

4. A compound of Claim 1 wherein R<sup>4</sup> is H.

5. A compound of Claim 1 wherein R<sup>7</sup> is H and R<sup>8</sup> is OH.

6. A compound of Claim 1 wherein R<sup>1</sup> and R<sup>5</sup> taken together form a group of the formula NH-Z-O, wherein Z is a linker comprised of from 1 to 6 carbon atoms and 0 to 2 nitrogen atoms and wherein the O is attached at the position of R<sup>5</sup>;

7. A pharmaceutical composition for use in the treatment of a disease or condition characterized by undesired cellular proliferation or hyperproliferation comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

8. A method for treatment of a disease or condition characterized by undesired cellular proliferation or hyperproliferation in a subject suffering therefrom, comprising administering a therapeutically effective amount of a composition of Claim 7 to the subject.

9. The method of Claim 8 wherein the disease is cancer.

10. A method for treatment of a disease or condition characterized by undesired cellular proliferation or hyperproliferation in a subject suffering therefrom, comprising the steps of:



- (a) administering to said subject a substantially sub-toxic dose of an Hsp90 client protein inhibitor;
- (b) waiting a period of time sufficient to allow development of a substantially efficacious response; and
- 5 (c) administering to said subject a synergistic dose of a benzoquinone ansamycin.

11. The method of Claim 10 wherein the Hsp90 client protein inhibitor is a protein kinase inhibitor.

10

12. The method of Claim 11 wherein the protein kinase inhibitor is selected from the group consisting of quinazoline, phenylamino-pyridine, pyrazolo-pyridine, pyrrolo-pyridine, indole, oxindole, benzyldene malononitrile, flavone, staurosporine, antibody, and ribozyme protein kinase inhibitors.

15

13. The method of Claim 11 wherein the protein kinase inhibitor is selected from the group consisting of N-(3-chloro-4-fluorophenyl)-7-methoxy-6-[3-(4-morpholinyl)propoxy]-4-quinazolinamine, 2-(2-chlorophenyl)-5,7-dihydroxy-8-[(3S,4R)-3-hydroxy-1-methyl-4-piperidinyl]-4H-1-benzopyran-4-one, 4-[(4-methyl-1-piperazinyl)methyl]-N-[4-methyl-3-[[4-(3-pyridinyl)-2-pyrimidinyl]amino]phenyl]benzamide, N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-Quinazolinamine monohydrochloride, and 3-[(3,5-dimethyl-1H-pyrrol-2-yl)methylene]-1,3-dihydro-2H-Indol-2-one.

20

14. The method of Claim 10 wherein the benzoquinone ansamycin is selected from the group consisting of a compound of Claim 1, 17-allylamino-17-desmethoxygeldanamycin, and 17-(2-(dimethylamino)ethylamino)-17-desmethoxygeldanamycin.

25

15. A method for treatment of a disease or condition characterized by undesired cellular proliferation or hyperproliferation in a subject suffering therefrom, comprising the steps of:

30

- (a) administering to said subject a synergistic dose of a benzoquinone ansamycin;
- (b) waiting a period of time sufficient to allow development of a substantially efficacious response; and
- 5 (c) administering to said subject a sub-toxic dose of an Hsp90 client protein inhibitor.


16. The method of Claim 15 wherein the Hsp90 client protein inhibitor is a microtubule stabilizing agent.

10 17. The method of Claim 15 wherein the microtubule stabilizing agent is selected from the group consisting of paclitaxel, an epothilone, discodermolide, and laulimalide.



15 18. The method of Claim 16 wherein the benzoquinone ansamycin is selected from the group consisting of a compound of Claim 1, 17-allylamino-17-desmethoxygeldanamycin, and 17-(2-(dimethylamino)ethylamino)-17-desmethoxygeldanamycin.


20 19. A method for treatment of a disease or condition characterized by undesired cellular proliferation or hyperproliferation in a subject suffering therefrom, comprising administering to said subject a composition of Claim 7 in combination with a cytotoxic drug selected from the group consisting of 5-fluorouracil, methotrexate, vinblastine, cyclophosphamide, mechlorethamine, chlorambucil, Melphalan, Ifosfamide, bleomycin, mitomycin and doxorubicin.

25 20. The method of Claim 10 or Claim 15 wherein the sub-toxic dose is between the maximum tolerated dose and one-twentieth of the maximum tolerated dose of the Hsp90 client protein inhibitor, and the synergistic dose is the maximum tolerated dose of the benzoquinone ansamycin.



geldanamycin

17-allylamino-17-desmethoxy-  
geldanamycin  
(17-AAG)17-(2-dimethylaminoethyl)amino-17-  
desmethoxygeldanamycin  
(17-DMAG)**FIGURE 1**



## FIGURE 2






FIGURE 3A




5

FIGURE 3B


10




Tarceva




Iressa




BIBX 1382




Gleevec




SU101




Lefunamide



Flavopiridol




CEP-701



UCN-01

**FIGURE 4**



5

FIGURE 5



Geldanamycin PKS Module 1 DNA Sequence: SEQ ID NO: 1

0001 TCCGACGAGC CGGTGGCGAT CGTGGCGATG GGCTGCCACC TGCCGGGCGA  
GGTCGCGACG  
5 0061 CCCGAGGACC TGTGGCGGTT GGTGGCCGAC GGGCGGGACG CGATCGCCGG  
GTTCCCGGAG  
0121 GACCGGGGCT GGGACCTGGC CGGGCTCTTC GACTCCGACC CGGACGCCGT  
GGGCAAGTCC  
0181 TATGTGCGCG AGGGCGGTTT CCTCACCGAC GCGGGCGGAT TCGACGCCGC  
10 ATTCTTCGGC  
0241 ATCTCGCCCC GTGAGGCGCT GGCGATGGAC CCGCAGCAGC GGTGCTGCT  
GGAGACCGCG  
0301 TGGGAGACCT TCGAGAATGC CGGAATCGAC CCGGGTTCGC TGCACGGCAC  
CGACGTCGGT  
15 0361 GTGTTTCAGCG GAGTGATGTA CCACGATTAC GGGGCCGACG CCGGGACGGC  
GGCGGAGGGC  
0421 CTGGAGGGGC ATCTCGGCGT GGGCAGCGCG GGGAGCGTCG TCTCCGGGCG  
GGTGGCCTAC  
0481 GCGCTGGGCC TGACCGGGCC CGCGGTGACC GTGGACACCG CCTGCTCGTC  
20 CTCCCTGGTA  
0541 GCGCTGCACC TGGCGGTTCA GGCGGTGCGC ACGGGCGAAT GCTCGCTGGC  
GCTCGCCGGG  
0601 GGTGTCGCGG TGATGAGCAG GCCGACGTCG TTCATCGAGT TCTCCCGCCA  
GCGTGGCCTC  
25 0661 GCCCCGACG GCCGCTGCAA GTCCTTCGCG GAGGGCGCCG ACGGCACCAA  
CTGGTCCGAG  
0721 GGTGTCGGGT TGGTGTGCT GGAGCGGCTG TCCGATGCCC GCCGCAATGG  
GCATGAGGTG  
0781 CTCGCCGTCG TCCGTGGCAC CGCCGTGAAC CAGGACGGCG CCAGCAACGG  
30 CCTGACCGCG  
0841 CCAACGGCC CGTCCCAGGA ACGGTGATC CGGCAGGCGC TGGCGAACGC  
CGGGCTGACG  
0901 GTGGCCGATG TGGACGCGGT CGAGGCCAC GGCACGGGCA CGAGTCTCGG  
CGACCCGATC  
35 0961 GAGGCCAGG CACTCCTGGC CACCTACGGG CAGGAGCGGC CGGAGGATCA  
GCCGCTGTGG  
1021 CTGGGGTTCGT TGAAGTCGAA CATCGGGCAT GCGCAGGCGG CGGCGGGCGC  
GGCCGGTGTG  
1081 ATCAAGATGG TCCAGGCCAT GCGGCACGGC GTACTGCCCA AAACCCTCCA  
40 CGCCGACGAG

1141 CCCACCAGCA AGGTCGACTG GACGTCAGGT GCGGTGTCGC TACTGTCCGA  
GGCCCCGGCCC  
1201 TGGCCGGAGA CGGGACACCC CCGCCGCGCC GGAATCTCCT CCTTCGGCGT  
CAGCGGGACG  
5 1261 AACGCACACG TGGTCCTGGA ACAGGCACCC CTGGAAGCGG CTGCACCCGA  
AACACAGGCG  
1321 AGCGACGCGG GCGCTCCTGG GCTCGTGGCC ACGGGCGGCG TAGTGCCGTG  
GGTGCTGTCC  
1381 GCCAAGACTC CTGCGGCGCT GCGCGCTCAG GCAGAGCGTC TGGTCAGCCA  
10 TCTGGAGTCC  
1441 GGGAGCGACG CCAACCCGGT CGATGTGGGC TGGTCGCTGG CCACCACCCG  
GGCGGCGTTG  
1501 GAGCACCGCG CGGTCATCCT GCGGACGGAT GCCGAAGGAG GCATGGCGAC  
GGCGCGGGCT  
15 1561 CTGGCGGAGG GCGGCCCTGA CCCGCTCCTG GTCACCGGAC AGACCGGAAC  
AGACGGCAAA  
1621 ACCGTGTTCA TCTTCCCCGG CCAAGGCGCC CAATGGGTGG GCATGGGAGC  
CCAACCTCCTC  
1681 AACACCTCAC CCGTCTTCGC CGCCCGCCTG CGTGAGTGCG CCGATGCTCT  
20 AGCGCCGTAT  
1741 ACCGACTGGT CGTCATCGA CGTCATCACC GGCACGCCCC ACGTCCCTC  
GCTTGAGCGT  
1801 GTCGACGTCG TACAGCCCGC CACCTTCGCC GTCGTGCTCT CCCTCGCCGC  
ACTCTGGCAA  
25 1861 TCCGTGGGCA TCCACCCCGA CGCCGTCATC GGCCACTCCC AAGGCGAAAT  
CGCCGCCGCC  
1921 TCGTGCGCCG GACACCTCAC CCTCACCAAC GCCGCCAAAA TCGTCACCCT  
CCGCAGCCAG  
1981 ACCATCGCCC ACCACCTCGC CGGACACGGC GGCATGATGT CCCTCGCCAC  
30 CCCC GCCGAC  
2041 ACCATCGACC TCACCAACTG GCACGGCAAA CTCTGGATCG CCGCACACAA  
CAGCCCCAAC  
2101 GCCACCGTCA TCGCAGGCGA CACCGACGCC CTGCACCAAC TCCACACCCA  
CTACACCGAC  
35 2161 CAGGGCACCA GAGCCCGCAT CATCCCCGTC GACTACGCCT CCCACACCGG  
ACACGTGAC  
2221 ACCATCAAAA ACCAGCTACA AGACGTACTC GACGGCATCA CCCTCGAACC  
CGGCACCATC  
2281 CCCTGGCTCT CCACCGTCAC CGGACAGTGG ATCGAACCCA ACACCGTCGG  
40 CGACAGCTAC

2341 TGGTACCGCA ACCTCCGCCA AACCGTGCAA TTCGAGCACA CCATCCACAC  
CCTCGCCGAC  
2401 CAGGGCTACC GCACCTACAT CGAAATCAGC CCCCACCCCG TCCTCACCAC  
CGCCATCCAA  
5 2461 GAAACCCTCG AAGCCAACGA CACCCCAAC ACCACCATCG TCACCGGCAC  
CCTCCGCCGC  
2521 GACGACGACA CCCCACCCG CCTCCTCACC AACCTCGCCC ACCTCACCAC  
CAACGGAACA  
2581 CCAGTCAACT GGCCACCCCT CTTACAGGC ACCCAACCCA CCCGCATCCC  
10 CCTCCCCACC  
2641 TACCCCTTCC AACACCACCA CTACTGGCTC CCCGCAACA CCAGCGCAGG  
CGATGTGAGT  
2701 GCCGTGGGCC TCCAGGGCAC GGGCCACCCG CTGGCCGGGG CCGTGGTGAG  
CGTGCCCGAC  
15 2761 ACCGGGGGTG TGCTGCTCAC CGGGCAGTTG TCGGTGGCCA CCCACCCGTG  
GCTGGCCGAC  
2821 CACGCCGTCT CCGGAACGGT GCTGCTGCCG GGCACCGCGA TGGCCGAAC  
CGCCATCCGC  
2881 GCCGGAGACG AGACCGACAC CCCCACCCTG GAAGAGCTGG TCATCGGCCA  
20 GCCGATGACA  
2941 CTGCCCCAAG ACGGTGCACT ACATGTCCAG GTACTGGTCG GCGGCGTGGA  
GGACGGGGCGC  
3001 CGAGGGGTGC GGATCTACTC TCGCCCCGAC GCGGCCCAGG AACAGGAATG  
GCTGGAGCAC  
25 3061 GCCTCGGGCA CACTCGCCAC GCAGCCGAC GGTTCGGCCG AGGGCGGCAT  
GGAGAACGGC  
3121 ATGCCCCAGT GGCCGCCGCC CGGTGTCGAG CCGATCGCTC TGGATGACTT  
CTACGACGAC  
3181 CTCGCCCAGG CCGGTATGA GTACGGGCCC GCCTTCCGCG GGCTGAAGGC  
30 GGTCTGGAAG  
3241 CGCGATGGCG AGGTGTTCGC GGAGGCCGCG CTGCCGAGG AGCAGACGGA  
CGTCGCCGGC  
3301 CGGTTCCGTA TCCATCCGGC GCTGCTGGAC GCCGCGTTGC ACGCGAGCAA  
CTTCTGTGTG  
35 3361 CCCCCGGCCC CGGGCCAAAC GCTCCTCCCC TTCGTGTGGA ACGGCGTACG  
GCTGCTGGCG  
3421 GCGGGAGCCA CGGCCGTCCG TGTGCGCGCC CGCGCCACCG GCACGGACTC  
GTTACGATC  
3481 AGCCTGTTCG ACAGCACCGG CTCCCCCGTC GCCTCGGTGG ACTCCCTGGT  
40 GCTCCGGGCG

3541 ATCAGTCCCG AGCAGCTCGC TGCCGCTCC GCGGTGCCG GTCGGTCCGC  
TGATGCGCTG  
3601 TTCACGCTGG ACTGGACCGA GCACCCACC GCCCTGGGGA CCGAGGTTTC  
CTGGGCCACC  
5 3661 CTCGGCGATG CCCACACCGA CGTGGACGCC CACGTGGACG CGCTCATCGC  
GGGAGAGGAC  
3721 CGGCCCCGGG CCGTGGTCGC CGACACCGCG GCCTGGGCCG CCGGGGACAC  
CGGCCTGCCC  
3781 GCGCGGGCCC GGGATCTGGC CGCCCGCGCG CTGGACCTGG TGCAGCGGTG  
10 GGTCGGCCGA  
3841 CCCGAAGTCG CCGACGTCCG GCTCGTGTTG CTCACTCGTG GGGCGGTGTC  
CGTGCACGAC  
3901 ACCGCCGAGG TCACCGACCC GGCCGCCGCC GCGATCTGGG GCCTGGTCCG  
CTCCGCCCAG  
15 3961 TCCGAACACC CGGGCCGGAT CGCCCTGGTG GACACCGACG ACGTGTCCG  
GGAGGCGCTG  
4021 CCCGAGGCGG TGGCGGCCGG CGAGCCGCAA GTGGCGCTGC GCCGTGGGCT  
GCTGTGGGTG  
4081 CCTCGTCTGG TCGGTTCGCC GCAGGGTCTC GCCGTACCCG CGCAGGAGCA  
20 CTGGTACCTC  
4141 GACGTCTCGG AGAAGGGCAG CCTGGAGAAC CTGGTGCTGC GGCCGGATCC  
GGAGGCCACC  
4201 GCGCCGCTGG CCACCGGTCA GGTCCGGATC GAGGTCCGCG CCGCCGGTCA  
GAACTTCCGG  
25 4261 GACGTACTCG TCGCGCTCGG CGGCGTGGCG GGTGAGGAGG GTCTGGGCGG  
CGAGGGTGCC  
4321 GGGGTGGTGA CCGAGGTCGG GCCCGGGGTC GAGGGCCTGG CGGTGGGCGA  
CCGGGTGATG  
4381 GGCCTGTTCC CGCGCTCGTT CGGCCCGCTG GCCATCGCGG ACGCGCGCAC  
30 GGTGCGGCCG  
4441 ATCCCGAGG GCTGGTCGTA CGCCACGGCC GCCGGGTGC CGGTGGCCTA  
TCTGACGGCA  
4501 CTGTACGGGC TCGGGACCT GGGCACCATA CAGCCGGGTG AGACGGTGCT  
GGTGACGCC  
35 4561 GCCGCGGGCG GTGTGGGCAT GGCCGCCGTC CAGTTGGCGC GGCCTTCGG  
CGCCACCGTG  
4621 TACGCCACCG CCCACCGTC GAAGCACCAT GTGCTGACCG CGCTGGGGGT  
GCCGGAGGGG  
4681 CATCTGGCGT CCAGCCGCGA CCTCGGTTTC GCCTCGGCGT TTCCCGCGCT  
40 GGATGTGGTG

4741 CTGAACTCCC TCACCGGCGA GTATGTGGAC GCCTCGCTGG GGCTGCTCGG  
CACGGGTGGC  
4701 CGTTTCGTGG AGATGGGCAA GAACGACATC CGCGATCCCG CCTCGGTGCG  
CGCAGCACAT  
5 4861 CCCGGTGTGG GCTATCAGGC GTTCGACCTG GGAGGTGACG CGGGCCCTGA  
CCGGATCCGG  
4921 GAGCTGCTCG CGGAGCTGGT GGAAGTGTTC GAGGCGGGCC GGATCGAGCC  
GCTTCCGATA  
4981 CGGCACTGGG ACGTCACCCA GGC GCCGACG GCCTTCCGGT GGATGAGCCA  
10 GGGGCGGCAC  
5041 ACCGGCAAGA TCGTGCTCAC CCTCCCCGA GCCCTGGACC CGGACGGCAC  
CGTCCTGATC  
5101 ACCGGTGGCA CCGGAACCCT CGGCGCCACC ATCGCCCGCC ACGTCGTCAC  
CCACCACGGC  
15 5161 GCGCGCCAGT TGCTCCTCAT CAGCCGTCAG GGTCCCGACG CCCCCGGCGC  
CACCGATCTC  
5221 ACCACCGAAC TCACCGAACT CGGCGCCACC GTCCGCATCA CCGCCTGCGA  
CACCGCCGAC  
5281 CGCGACCAAC TCGCCGCGCT CCTCGCCGAC ATCCCCGCCG CCCACCCCT  
20 CACCGCCGTC  
5341 ATCCACACCG CCGGCGCCCT GGACGACGGT GTCCTGACCG CGCTACCCCC  
GGACCGCCTC  
5401 GACACCGTCT TCCGCCCAA GGTCGACGCC GTCACCCACC TCCACGACCT  
CACCCGCGAC  
25 5461 CAGGACCTGG CCGCGTTCGT CATCTACTCG TCCGCCGCCG GAACGCTCGG  
CAACGCGGGG  
5521 CAGGCCAACT ACGCCGCCGC CAATGCCTTC CTCGACGCCT TCGCCAGTG  
GCGGCACGCC  
5581 CGCCACCGGC CCGCCACTTC GCTGGCGTGG GGGCTGTGGA GCGACACCAG  
30 CACGCTCACC  
5641 TCGACGATGG ACGCCACCGA CGTACGCCGC ACACGGCGGG CGGGGGTGCT  
GGGCATGGAC  
5701 AACGCCGAGG CGTGCGGGT GTTCGACACC GGGTTGCGGT CCGGGCGGCC  
CGCGCTGGTG  
35 5761 GCCGCGAAGA TCGACCTCAC CGCCCTGCGC GCGCCGACG CCGAGTTGTC  
GCCGCTGCTG  
5821 CGCGGACTGG CCCGTCCGGC GCGCCGACCG GCGCGACCG CGGCCCCGGC  
GGCCGGTGGT  
5881 CTGTCGGGGC AGCTGGCCGG GCTGTCCCCC GCCGGGCAGC GGGAGTTCCT  
40 GCTCAACCTG

5941 GTGCGGGCGG AGGCCGCGGT GGTCTCTGCC CACGCCGGTC CTGAGGCGAT  
CGAGCCGACC

6001 GTGGCGTTCA AGGAGATGGG TTTCGACTCG CTGACGGCGG TCGAACTGCG  
CAACCGGCTG

5 6061 AATGCGGCGA CCGGGCTGCG GCTCCCCGCC ACGTTGCTCT TCGACCACCC  
GACTCCGGCT

6121 CTTCTACCG AGCTGTTCCA TACCGAGTTG GCGGGCGGCC CGGCACCCGC  
CGCGGCGGCC

6181 CCGGTGACCG TCGTGCCGC CGCTGACGAG //

10

Geldanamycin PKS Module 1 DNA Sequence: SEQ ID NO: 2

0001 ATGGCTGCGT CCCGGAAGA CCTGGTCAAG GCGCTGCGTA CCTCGCTGAT  
GGACGCCGAG  
5 0061 CGGCTGAAGC GGGAGAACGA CCGGCTGATC GCCGAGTCCA CCGAACCGGT  
GGCGATCGTG  
0121 GCGATGGCGT GCCGGCTGCC GGGTGGGGTG ACCGACCCGG AGTCGCTGTG  
GGAGCTGGTG  
0181 GACGAGGGGC GGGACGCGAT CGGGCCGTTT CCCACGGATC GCGGCTGGGA  
10 CCTGGAGACC  
0241 CTGTTTCACT CCGATCCGGA CGCCGTGGGC AAGTCCTACG TACGCGAGGC  
GGGGTTCCTG  
0301 GAGGGGGCGG GCGGATTCGA CGCCGCCTTC TTCGGCATCT CGCCGCGCGA  
GGCCCTGTCTG  
15 0361 CTGGACCCGC AGCAGCGGCT GCTGCTGGAG ACCGCGTGGG AGACCTTCGA  
GCGGGCGGGG  
0421 ATGGATCCGC GGTCCGTGGA GGGCCGGGAC ATCGCGGTGT TCGCCGGGGG  
CAGCGGCCAG  
0481 GGGTACGGCG GCGGTCCGGG TGAGGCGCCC AAGGGCCTGG AGGGCTATCT  
20 GGGGGTCGGC  
0541 GCTTCCGGCA GTGTATCTC CGGGCGCGTG TCGTACACGC TCGGGCTGAC  
CGGTCCCGCC  
0601 GTGACCGTGG ACACCGCCTG CTCGTCTCG CTGGTGGCCG CCCATCTCGC  
CGTGCAAGCG  
25 0661 CTGCGGTCCG GCGAATGTTC CATGGCGCTG GCCGGTGGTG TCGCCGTGAT  
GGGCCAGCCC  
0721 ACCGCCTTCG TCGAGTTCTC CCGGCAGCGT GGCCTGGCGC CCGACGGGCG  
CTGCAAGTCC  
0781 TTCGGCGCGG GCGCCGACGG CACCACCTGG TCCGAAGGTG TCGGGCTCGT  
30 TCTGCTGGAG  
0841 CGGCTGTCGG ACGCCCGCCG CAACGGCCAC GAAGTGCTGG CCGTGATCCG  
GGGCACCGCG  
0901 GTCAACCAGG ACGGCGCCTC CAACGGACTC ACCGCGCCCA ACGGCCCTC  
CCAGGAGCGG  
35 0961 GTGATCCGCC AGGCCCTGTC CAACGCCGGG CTGACGGTGG CCGACGTGGA  
CGCCGTCGAG  
1021 GCCACGGCA CCGGCACCGC CCTCGGCGAC CCCATCGAAG CCCAGGCCGT  
TCTCGCCACC  
1081 TACGGCCAAA GCCGCCCGGA GGGCCGGCCG CTGTGGCTCG GCTCCCTCAA  
40 GTCCAACATC

1141 GGCCACGCGC AGGCCGCAGC GGGCATCGCC AGTGTCATCA AGACCGTCAT  
GGCCTTACGC  
1201 CACGGCCGGT TGCCGAAGAC CCTCCACGCC GAACAGCCCA CCTCCCAGGT  
GAACTGGACG  
5 1261 TCGGGCGCGG TGTCCCTGCT CGCCGAGGCG CGGGCGTGGC CGGAGACCGG  
ACACGCCCCG  
1321 CGCGCCGGGA TCTCCTCCTT CGGCGTCAGC GGGACGAACG CACACGTCAT  
CCTGGAACAG  
1381 GCCCCTGAGG AAGCCGAGGC GACCGGGGAG AACACCGCCG ATCAGGAACC  
10 GCCCGTACGC  
1441 TCGGCGGAGT CCGCCGACCC CGGCCCGGTC GCCACCGGCC ACGTGGTGCC  
GTGGCTGCTC  
1501 TCGGGCCATA CGCAGGAGGC GCTGCGTGCC CAGGCCGCCG GGCTGCTGAC  
CCAGGTGCGC  
15 1561 GAGACGCCCT CCGACAGTCC GCGGGACGTG GGCTGGTCAC TGGCCACCAC  
CCGGACCCGG  
1621 CTGGACCACC GCGCGGTCGT ACTGTGCGCC GATGCCGAGC AGGCCGTCGC  
GGGGCTGGAG  
1681 GCGGTGGCCT CGGGCACGTC CGCCCGGTG GCGGTCACCG GGTCCGTGGC  
20 CTCCGGAAAG  
1741 GTGGCGGTGC TGTTCACCGG GCAGGGCAGC CAGCGGGCCG GAATGGGCCG  
CGAACTGCAC  
1801 GGCGCCCACC CGGTGTTGCG GCGGGCCTTC GACGCCGTGT GCGCCCAGTT  
CGGCGACCTG  
25 1861 CGCGACGGG ACGACAAGGT CTCGCTGGCC GAGGTGATCT TCGCCGAGGA  
GGGGTCGGCG  
1921 ACGGCAGCGC TGCTGGACCG GACCGAGTTC ACCCAGCCCG CGCTGTTGCG  
GCTGGAGGTG  
1981 GCGCTGTTCC GGCTCGTGGA GTCGTGGGGA GTGCGCCCCG CGTATGTGCT  
30 GGGCCACTCG  
2041 ATCGGCCAAG TGGCGGCGGC CCATGTGGCC GGGGTCCTGT CCCTGCCGGA  
CGCCTGCACA  
2101 TTGGTGCGGG CGCGCGGGCG GCTGATGCAG CAACTCACCG CGACCGGGGC  
GATGGTCGCG  
35 2161 GTGAGGCGG CCGAGGACGA GGTGGCGCCG CTGCTCGCGG GGAAGGAGCA  
CAAGGTCTCC  
2221 ATCGCCGCGG TCAACGGCCC GGCCTCCGTG GTCGTCTCCG GTGACGAGGA  
CGTGGTCACG  
2281 GCGGTGGCGG AGACGCTGGC GCGGCAGGGC CGCAAGACCA AGCGGCTCGT  
40 GGTCTCGCAC



2341 GCCTTCCACT CCCCCACAT GGACGGGATG CTGGACGCGT TCCGCGAGGT  
GGCGTCGCGG  
2401 CTGGCCTACG CGCCACCCCG GATACCCGTG GTGTCGAACC TCACCGGCGC  
GGTCGCCGAT  
5 2461 CCCGAGGAGC TGTGCTCCCC CGAGTACTGG GTACGGCATG CACGTGGCGC  
GGTGCGGTTC  
2521 CTCGACGGTG TCCGCACACT GGCEGACGAG GGCCTGCGCA CCCATCTGGA  
ACTCGGCCCC  
2581 GATGGGGTGC TGACCGCGAT GGGGACAGGAC TGTCTGCCCC AGGCGGACGC  
10 GCGTTTCGTG  
2641 CCGTCCCTGC GTCCGGGCGT CCAGGAGCCG CACGCGGTGC TGGCCGGGCT  
CGCCGGCCTG  
2701 TACGTACGGG GTGTGCGGGT GGA CTGGGAC GCGATGTTCTG CCGGGTCCGG  
CGCCCGGCCC  
15 2761 GTCGCCCTTC CCACGTACGC CTTCCAGCAC GAGCACTACT GGCTGGAGCG  
GGCCGCCGGC  
2821 TCCGGCGACG TGGGCGCGGT GGGGCTCGGC GAGGCGGGCC ATCCGCTGCT  
GGGCGCGGTG  
2881 GTGCAGCTCC CGGAGACGGG CGGGGTGCAG CTCAGCGGGC GGCTGTGCGT  
20 ACGGGCCCCAG  
2941 CCCTGGCTGG GCGAACACGT CATCTCCGGG GCGGTGCTGG TGCCCGGCAC  
CGCCATGGTG  
3001 GAACTGGCCG TCCGCGCCGG GGACGAGACC GGCACCCCGG TGCTGGAGGA  
GCTGGTGATC  
25 3061 GGGCAGCCGA TGGTGTGCC CGGCGACACC GCCCTCAGTG TCCAGGTCGT  
CGTGGGCGCG  
3121 GACGAGGGCG GCGGCGGTAC GGTGCGGATC TACTCCCGTA CCGACGGGGG  
CACCGACTGG  
3181 ACCGAGCACG CCACCGGCAC GCTCGCGGCG CAGGGCCCGG CACCGCTGGA  
30 CGGGGCCCGG  
3241 GCGGGGGCCG CCGTCGAGTG GCCGCCCGCG GAAGCCGAGC CGATCCCCGT  
GGAGGACTTC  
3301 TACCGCTCGC TCGTCGACGC CGGATACGCG TACGGACCGG CGTTCCGCGG  
GCTCGTCGCC  
35 3361 GCGTGGCGCC GGGACGGTGA GATCTTCGGC GATGTGGCGC TGCCGGAGGC  
GTCCGTGCGG  
3421 GAGGCCGAGC GGTTCGGCAT CCACCCGGCG CTGCTGGACG CCGCACTGCA  
CGCGGGCAGC  
3481 TTCTGTCTGC CCTCCGACCC GCGCGACAG GTGACCCTGC TGCCGTTCGC  
40 CTGGAACACC

3541 GTGCGTCTGC ACGCGGGCGG CGCGTCCGCG GTCCGGGTGC ATGTCCGCCC  
GGTCGGCGAC  
3601 GACGCCTTCT CGGTACGCCT GACCGACGGC TCGGGCCAGA CGGTGGCCTC  
GGTGGACTCG  
5 3661 CTCACCTTGC GGGCGGTGGA CCCGGCCCAG CTCAAGATCG GCACGGCCGA  
CGACGCGCTG  
3721 TGGACGGTCC GCTGGAGCGA GACCTCGCTG CCGACGGCG CGGTCTCCTG  
GGCCCCGCTC  
3781 GGCAGATCGG CCACCGGGG AACCAGGGG TACGGCGCCA CAGGGGACGG  
10 CGGAGGCCCA  
3841 GGGGGCGCGC TTCCCGACGT CCTCGTGGCC GATACGCGCG CCTGGGCCGA  
AGACCTCACC  
3901 GGACCGCCGA CCGCGCGGGC CCGGAGCTC ACCGGCCGCC TGCTGGAGGA  
GATCCAGCGG  
15 3961 TGGGTCGCCG ACGACCCAT GGCCGGGACG CGGCTCGCCG TGGTCACCCG  
CGGCGCGGTC  
4021 GCGGTCCACG ACGACACCGA GGTACCGAC CCGGCCGCA CCGCGCTCTG  
GGGCCTGGTC  
4081 CGCTCGGCCC AGGCCGAACA CCCGGGGCGG GTGGCCCTGG TGGATGCCGA  
20 CGGAGCGTGC  
4141 GAGGAACTGC CCGCCGGGGT GTGGTCCGGG GACGAGCCCC AACTGGCGGT  
GCGCGGTGGC  
4201 GCCGTGTGGG TGCCACGCCT CACCCGGGTC GAGCCCGGCC TCGCGTGCC  
CGCGCAGGCG  
25 4261 TCGTGGCATC TGGACTCGGC CGAGTACGGC ACCCTGGACA ATCTGGCGCT  
GCTGCCCGAC  
4321 GAGGCCGAGC CCGCACCGCC GGCGGCCGGT CAGGTGCGGA TCGAGGTCCG  
CGCCGCCGGG  
4381 CTCAACTTCC GGGATGTCTT GGTGGCTCTC GGCATGTATC CGGGCCGGTC  
30 GGTGATCGGC  
4441 ACGGAGGGCG CCGGTGTGGT GACCGAAGTC GGTCCGGGCG TCACGGGCCT  
GGCCGTGGGC  
4501 GACCGGGTGA TGGCCTGTT CTCCGGCTCG TTCGGACCGC TGGCCACCGC  
CGACGCGCGC  
35 4561 ACGGTGATCC GGATGCCGGA GGGCTGGTCG TTCGGCACGG CGGCCGGGT  
GCCGGTGGCC  
4621 TATCTGACGG CGCTGTACGC GTTGCAGGAC CTCGGGAGGG TCCAGCCGGG  
CGAGACGGTC  
4681 CTGGTGCACG CCGCCGCGGG CGGTGTGGGC ATGGCCGCCG TCCAGCTCGC  
40 ACAGCACTTC

4741 GGC GCCACCG TCCTGGGCAC CGCCACCCC TCCAAGCACC ACGCACTCCA  
CCGGCTGGGC  
4801 GTTCCCGCCG AACGGCTCGC CTCCAGCCGC GACCTCGCCT ACGCCGACAC  
CTTCCCCACC  
5 4861 GCCGACGTCG TCCTCAACTC CCTCACC GGC GAGCACATCG ACGCCTCCCT  
CGGACTTCTC  
4921 AACCCCGGCG GCCGGTTCCT GGAGATGGGG AAGACCGACC TGC GGGAGCC  
CGGCGAGGTC  
4981 GGGCGCGGC ATCCGGAGGT CACCTACCGG GCGTTCGATC TCGGTGGGGA  
10 GGCCCCCGC  
5041 GAGCGGTGCG GGGAGTTGCT GCACCAGTTG GTGGAGCTGT TCGAGGCGGG  
CCGGATCGAG  
5101 CCGTGCCCG TACGGCAGTG GGACATCACC CGCGCCCCCG AGGCGTTCCG  
CTGGATGAGT  
15 5161 CAGGGCGGC ATACCGGCAA GATCGTGCTC ACCCTGCCAC GCGCCCTGGA  
CCCGGACGGC  
5221 ACCGTCCTGG TCACCGGTGG CACGGGCACC CTCGGCGCCA CGATCGCCCC  
CCACCTTCTC  
5281 ACCCAGCAG GCGCACGCCA TCTGCTGCTG GTCAGCCGCC GGGGACCGGA  
20 CGCACCTGGC  
5341 GCCACAGACC TGACCACCGA ACTCACC GAA CTCGGCGCCA CCGTCCGCAT  
CACCGCCTGC  
5401 GACACCGCCG ACCGCGACCA ACTCGCCGCG CTCCTCGCCG ACATCCCCCG  
CGACCACCCC  
25 5461 CTCACCGCCG TGGTCCACAC GGCCGGGACC CTCGACGACG GTGTCCTGAC  
CGCGCTCACC  
5521 CCGGACCGCC TCGACACCGT CTTCCGCCCC AAGGTCGACG CCGTCACCCA  
TCTCCACGAC  
5581 CTCACCCGCG ACCACGACCT GGC GCGTTC GTGGTGTACT CGTCCGCCG  
30 CGGAGTCCTC  
5641 GCGGGGCCCC GCCAGGGCAA CTACTCCGCC GCCAACGCCT ATCTGGACGG  
ACTCGCACAG  
5701 TGGCGGCGTG CGCACGGGCT CCCC GCCACC TCGCTGGCGT GGGGCATGTG  
GGCGCAGACC  
35 5761 AGTGGCATGA CGGCCGGGCT CGGCTCCGGC GATCTGCACC GGGTCCGGCG  
TGGCGGCATC  
5821 GTCGGGCTGT CCACGGCGGA GGCCCTGGAC CTGTTCGACC GGTCCGTGGC  
GTCCGGGCTG  
5881 TCCCTGCTGG TGCCGTTGCG GTTGACATC GCCGCCCTCG GTGCGGAGGC  
40 CGCGGAACCG

5941 CCGCCGCTGC TCGGGGTCT GGTCCGGCCG GCCCGGCGTA CGGCCCGGCC  
GGTGCCGAAG  
6001 GCCGGTGAGG GCGGCCTCGC CGAACGGCTG GCCGGGCTGT CGGCGGCCGA  
ACAGGAGCGT  
5 6061 CTGCTCATCG AGTTGATCCG CGAACAGGCC GCTTCGGTGC TCGGGTCCC  
CACGGTCGAC  
6121 CCGATCGGGC CGGAGCAGGC GTTCCGCGAC ATGGGGTTCG ACTCGCTGAC  
CGCGGTGGAG  
6181 CTGCGCAACC GCCTCAACAC GGCCACCGGG CTACGGCTCC CCGCAACGCT  
10 GGTCTTCGAC  
6241 CACCCGAGCC CCTTGGCCAC CGCCGAGTTC CTGCGGGATC AACTGGGCGG  
GCGCGCGGTC  
6301 GAGGCGGCGC CCCGCCGGC CCGGCGTGAC CGGTCGGCTC CGGACGGGGC  
CGAGGATCCG //  
15

20 Specificity Peptide Sequence of module 1 AT domain: SEQ ID NO: 3

0001 Tyr Ala Ser His //

Specificity Nucleotide Sequence of module 1 AT domain: SEQ ID NO: 4

25 0001 tac gcc tcc cac //

Specificity Peptide Sequence of mutated module 1 AT domain: SEQ ID NO: 5

0001 His Ala Phe His //

30 Specificity Nucleotide Sequence of module 1 AT domain: SEQ ID NO: 6

0001 cac gcc ttc cac //

Signature Peptide Sequence of module 6 DH domain: SEQ ID NO: 7

35 0001 His Val Ile Ser Gly Ala Val Leu Val Pro //

Signature Nucleotide Sequence of module 6 DH domain: SEQ ID NO: 8

0001 cac gtc atc tcc ggg gcg gtg ctg gtg ccc //

Signature Peptide Sequence of module 6 DH domain: SEQ ID NO: 9

5

0001 His Ala Val Ser Gly Thr Val Leu Leu Pro //

Signature Nucleotide Sequence of module 6 DH domain: SEQ ID NO: 10

0001 cac gcc gtc tcc gga acg gtg ctg ctg ccg //

10

